Biochimie 81 (1999) 1133–1138
© Société française de biochimie et biologie moléculaire / Éditions scientifiques et médicales Elsevier SAS. All rights reserved.

Author index 1999

A

Aarsman M.E.G. – See Roos M. et al., 797
Accorsi A. – See Piatti E. et al., 1011
Acebrón A. – See Ros P. et al., 389
Acquaviva R. – See Pacifico F. et al., 493
Ahn J.Y., Choi S., Cho S.W. – Identification of lysine residue involved in inactivation of brain glutamate dehydrogenase isoproteins by o-phthalaldehyde, 1123
Aledo J.C., Segura J.A., Barbero L.G., Márquez J. – Upregulation of glyceraldehyde-3-phosphate dehydrogenase mRNA in the spleen of tumor-bearing mice, 1109
Alex S. – See Colzani R.M. et al., 485
Alexandre S. – See Norris V. et al., 915
Alexciev K. – See Escobar Galvis M.L. et al., 1089
Alix A.J.P. – See Debelle L. and Alix A.J.P., 981
Almouzni G. – See Moggs J.G. and Almouzni G., 45
Aloj S.M. – See Bifulco M. et al., 287
An W., Zlatanova J., Leuba S.H., van Holde K. – The site of binding of linker histone to the nucleosome does not depend upon the amino termini of core histones, 727
Anagnostides S.T. – See Chatziioannidis C.C. et al., 187
Aoki H. – See Kiel M.C. et al., 1097
Arai K.M. – See Masai H. et al., 847
Arluison V., Batelier G., Riès-Kautt M., Grosjean H. – RNA:pseudouridine synthetase Pus1 from *Saccharomyces cerevisiae*: Oligomerization property and stoichiometry of the complex with yeast $tRNA^{Phe}$, 751
Ayala J.A. – See Carrión M. et al., 879

B

Bandorowicz-Pikula J. – See Danieluk M. et al., 717
Barbero L.G. – See Aledo J.C. et al., 1109
Barbier-Brygoo H., Joyard J. – Functions and dynamics of plant membranes, 575
Batelier G. – See Arluison V. et al., 751
Bechtner G., Fröschl H., Sachse A., Schopohl D., Gärtner R. – Induction of apoptosis in porcine thyroid follicles by transforming growth factor β1 and epidermal growth factor, 315
Beckers C. – See Van Den Hove M.F. et al., 563
Bedouelle H. – See Jermutus L. et al., 235
Behrends J. – See Pirson I. et al., 309
Belfiore A., Pandini G., Vella V., Squatrito S., Vigneri R. – Insulin/IGF-I hybrid receptors play a major role in IGF-I signaling in thyroid cancer, 403
Belisario M.A. – See Pelagalli A. et al., 1003
Bensimon A. – See Herrick J. and Bensimon A., 859
Benvenga S., Robbins J. – Altered thyroid hormone binding to plasma lipoproteins in the syndrome of resistance to thyroid hormones, 545
Benvenuto G. – See Chiariotti L. et al., 381
Bergmann E. – See Frit P. et al., 27
Bermúdez-Cruz R., Chamberlin M.J., Montañez C. – Nus A is involved in transcriptional termination on λ *tI*, 757
Biagini A. – See Giardina T. et al., 1049
Bianchi N. – See Mischiati C. et al., 219
Bifulco M., Laezza C., Aloj S.M. – Inhibition of farnesylation blocks growth but not differentiation in FRTL-5 thyroid cells, 287
Bigler S. – See Gerber H. et al., 441
Binenbaum Z., Klyman E., Fishov I. – Division-associated changes in membrane viscosity of *Escherichia coli*, 921
Bineta M.N. – See Lebrun-Garcia A. et al., 663
Bisi H. – See Tomimori E.K. et al., 447
Biville F. – See Perrotte-Piquemal M. et al., 245
Blaesing F. – See Messer W. et al., 819
Bliznakov E.G. – Aging, mitochondria, and coenzyme Q_{10}: The neglected relationship, 1131
Block M.A. – See Sato N. et al., 619
Boer J. de, Hoeijmakers J.H.J. – Cancer from the outside, aging from the inside: Mouse models to study the consequences of defective nucleotide excision repair, 127
Boevink P. – See Satiat-Jeunemaitre B. et al., 597
Bohr V.A., Dianov G.L. – Oxidative DNA damage processing in nuclear and mitochondrial DNA, 155
Boiteux S., Radicella J.P. – Base excision repair of 8-hydroxyguanine protects DNA from endogenous oxidative stress, 59
Bono J.J. – See Niebel A. et al., 669
Bouché J.P. – See Dassain M. et al., 889
Boudou V. – See Gaubert G. et al., 1041
Bouligand Y. – See Norris V. et al., 915
Bourque S. – See Lebrun-Garcia A. et al., 663
Braverman L.E. – See Colzani R.M. et al., 485
Bravo C., Woisard A., Fourrey J.L., Laugâa P., Favre A. – A Y form of hammerhead ribozyme trapped by photo-cross-links retains full cleavage activity, 201
Brino L., Bronner C., Oudet P., Mousli M. – Isoleucine 10 is essential for DNA gyrase B function in *Escherichia coli*, 973
Bronner C. – See Brino L. et al., 973
Bruni C.B. – See Chiariotti L. et al., 381
Bucchini A. – See Piatti E. et al., 1011
Bürgi U. – See Gerber H. et al., 441
Burrow G.N. – See Eggo M.C. et al., 291
Busch K. – See Fujiwara T. et al., 213

C

Calsou P. – See Salles B. et al., 53 – See Muller C. et al., 117
Camargo R.Y.A. – See Tomimori E.K. et al., 447
Cammenga M. – See Graaf S.A.R. (van de), 425
Campbell A.K. – See Holland I.B. et al., 901
Carlomagno F. – See Santoro M. et al., 397
Carolé S. – See Dassain M. et al., 889
Carr A.M. – See Caspari T. and Carr A.M., 173
Carrión M., Gómez M.J., Merchante-Schubert R., Dongarrá S., Ayala J.A. – *mraW*, an essential gene at the *dcw* cluster of *Escherichia coli* codes for a cytoplasmic protein with methyltransferase activity, 879
Carvalho D.P. – See Leseney A.M. et al., 373
Casamassima A. – See Vitale M. et al., 477
Caspari T., Carr A.M. – DNA structure checkpoint pathways in *Schizosaccharomyces pombe*, 173
Cattaneo A. – See Ojha M. et al., 765
Cavarelli J. – See Martinis S.A. et al., 683
Cellier D. – See Norris V. et al., 915
Cellini E. – See Degl'Innocenti D. et al., 1031
Chamberlin M.J. – See Bermúdez-Cruz R. et al., 757
Chanson P. – See Leseney A.M. et al., 373
Chany C. – See Jiang P.H. et al., 701 – See Kaba A. et al., 709
Chany-Fournier F. – See Jiang P.H. et al., 701 – See Kaba A. et al., 709
Chatziioannidis C.C., Karamanos N.K., Anagnostides S.T., Tsegenidis T. – Purification and characterisation of a minor low-sulphated dermatan sulphate-proteoglycan from ray skin, 187
Chiariotti L., Salvatore P., Benvenuto G., Bruni C.B. – Control of galectin gene expression, 381
Chieffi P., Picascia A., Stanzione R., Villone G., Tramontano D. – Endogenous insulin-like growth factors regulate the proliferation of TSH-independent mutants derived from FRTL5 cells, 367
Chiltz A. – See Lebrun-Garcia A. et al., 663
Cho S.W. – See Ahn J.Y. et al., 1123
Choi S. – See Ahn J.Y. et al., 1123
Churchich J.E. – See Kwon O.S. and Churchich J.E., 1057
Clark B.F.C. – See Wind T. et al., 1079
Cock J.M. – See Giranton J.L. et al., 675
Colosetti L. – See Dassain M. et al., 889
Colzani R.M., Alex S., Fang S.L., Stone S., Braverman L.E. – Effects of iodine repletion on thyroid morphology in iodine and/or selenium deficient rat term fetuses, pups and mothers, 485
Combettes B., Grienenberger J.M. – Analysis of wheat mitochondrial complex I purified by a one-step immunoaffinity chromatography, 645
Consiglio E. – See Suzuki K. et al., 329 – See Pacifico F. et al., 493 – See Di Jeso B. et al., 497
Corda D. – See Di Cerbo A. and Corda D., 415

Couprie J. – See Scala D. et al., 771
Cozzone A.J. – See Prost J.F. and Cozzone A.J., 197
Cretin F. – See Tapparo A. et al., 943
Cullimore J. – See Niebel A. et al., 669
Curcio F. – See Meli A. et al., 281

D

d'Alençon E., Taghbalout A., Kern R., Kohiyama M. – Replication cycle dependent association of SeqA to the outer membrane fraction of *E. coli*, 841
d'Angelo D. – See Pelagalli A. et al., 1003
Danchin A. – See Perrotte-Piquemal M. et al., 245
Danieluk M., Pikula S., Bandorowicz-Pikula J. – Annexin VI interacts with adenine nucleotides and their analogs, 717
Dantzer F., Schreiber V., Niedergang C., Trucco C., Flatter E., De La Rubia G., Oliver J., Rolli V., Ménissier de Murcia J., de Murcia G. – Involvement of poly(ADP-ribose) polymerase in base excision repair, 69
Dapas B. – See Morassutti C. et al., 1115
Das R., Vothknecht U.C. – Phenylalanyl-tRNA synthetase from the archaeon *Methanobacterium thermoautotrophicum* is an $(\alpha\beta)_2$ heterotetrameric protein, 1037
Dassain M., Leroy A., Colosetti L., Carolé S., Bouché J.P. – A new essential gene of the 'minimal genome' affecting cell division, 889
De La Rubia G. – See Dantzer F. et al., 69
de Murcia G. – See Dantzer F. et al., 69
De Nayer P. – See Van Den Hove M.F. et al., 563
De Riu S. – See Vitale M. et al., 477
De Vita G. – See Santoro M. et al., 397
de Vitry C., Vallon O. – Mutants of *Chlamydomonas*: Tools to study thylakoid membrane structure, function and biogenesis, 631
De Zegher F. – See Van Den Hove M.F. et al., 563
Debelle L., Alix A.J.P. – The structures of elastins and their function, 981
Degl'Innocenti D., Marzocchini R., Rosati F., Cellini E., Raugei G., Ramponi G. – Acylphosphatase expression during macrophage differentiation and activation of U-937 cell line, 1031 – ***, Rosati F., Iantomasi T., Vincenzini M.T., Ramponi G. – GSH system in relation to redox state in dystrophic skin fibroblasts, 1025
DeGroot L.J. – See Takeda T. et al., 297
Delamarche C., Guerdoux-Jamet P., Gras R., Nicolas J. – A symbolic-numeric approach to find patterns in genomes. Application to the translation initiation sites of *E. coli*, 1065
Della Morte R. – See Pelagalli A. et al., 1003
Demarty M. – See Norris V. et al., 915
Dème D. – See Leseney A.M. et al., 373
Deneke J. – See Masai H. et al., 847
Desideri A. – See Santucci R. et al., 745
Devlieger H. – See Van Den Hove M.F. et al., 563
Di Cerbo A., Corda D. – Signaling pathways involved in thyroid hyperfunction and growth in Graves' disease, 415
Di Jeso B., Formisano S., Consiglio E. – Depletion of divalent cations within the secretory pathway inhibits the terminal glycosylation of complex carbohydrates of thyroglobulin, 497
Di Lauro R. – See Macchia P.E. et al., 433
Dianov G.L. – See Bohr V.A. and Dianov G.L., 155
Dongarrá S. – See Carrión M. et al., 879
Doumas P. – See Santonia V. et al., 655
Dumas C. – See Giranton J.L. et al., 675
Dumont J.E. – See Pirson I. et al., 309
Dunn A.D. – See Dunn J.T. and Dunn A.D., 505
Dunn J.T., Dunn A.D. – The importance of thyroglobulin structure for thyroid hormone biosynthesis, 505
Dupuy C. – See Leseney A.M. et al., 373

E

Eckardt-Schupp F., Klaus C. – Radiation inducible DNA repair processes in eukaryotes, 161
Eggo M.C., Vitale M.A., Petric M., Burrow G.N. – Effects of thyroid-stimulating hormone on the replicative cycle of vesicular stomatitis virus in primary cultured sheep thyroid cells, 291
Egly J.-M. – See Frit P. et al., 27
Ehrenberg M. – See Karlsson M. et al., 995
Einav M. – See Zaritsky A. et al., 897
Eriksson S. – See Gaubert G. et al., 1041
Escargueil A.E. – See Scala D. et al., 771
Escobar Del Rey F. – See Escobar-Morreale H.F. et al., 453
Escobar Galvis M.L., Håkansson G., Alexciev K., Knorpp C. – Cloning and characterisation of a pea mitochondrial NDPK, 1089
Escobar-Morreale H.F., Obregón M.J., Escobar Del Rey F., Morreale De Escobar G. – Tissue-specific patterns of changes in 3,5,3´-triiodo-L-thyronine concentrations in thyroidectomized rats infused with increasing doses of the hormone. Which are the regulatory mechanisms?, 453
Escrivá H., Rodríguez-Peña A., Vallejo C.G. – Expression of mitochondrial genes and of the transcription factors involved in the biogenesis of mitochondria Tfam, NRF-1 and NRF-2, in rat liver, testis and brain, 965

F

Fairbank A.T. – See Hilderman R.H. and Fairbank A.T., 255
Fang S.L. – See Colzani R.M. et al., 485
Favre A. – See Bravo C. et al., 201
Faye L. – See Gomord V. et al., 607
Feldt-Rasmussen U. – See Rasmussen A.K. et al., 511
Fenzi G. – See Macchia P.E. et al., 433
Fenzi G.F. – See Vktdlg M. et al., 477
Feriotto G. – See Mischiati C. et al., 219
Fishov I. – See Zaritsky A. et al., 897 – See Binenbaum Z. et al., 921
Flatter E. – See Dantzer F. et al., 69
Florentz C. – See Martinis S.A. et al., 683
Fontes R., Günther Sillero M.A., Sillero A. – Acyl-CoA synthetase catalyzes the synthesis of diadenosine hexaphosphate (Ap_6A), 229
Formisano S. – See Suzuki K. et al., 329 – See Pacifico F. et al., 493 – See Di Jeso B. et al., 497
Fourrey J.L. – See Bravo C. et al., 201
Fraternale D. – See Piatti E. et al., 1011
Friedberg E.C. – The discovery of enzymatic photoreactivation and the question of priority: The letters of Salvador Luria and Albert Kelner, 7
Frit P., Bergmann E., Egly J.-M. – Transcription factor IIH: A key player in the cellular response to DNA damage, 27 – See Muller C. et al., 117
Fröschl H. – See Bechtner G. et al., 315
Fujiwara T., Busch K., Gross H.J., Mizutani T. – A SECIS binding protein (SBP) is distinct from selenocystyl-tRNA protecting factor (SePF), 213
Furui Y. – See Masai H. et al., 847
Fusco A. – See Santoro M. et al., 397

G

Gambari R. – See Mischiati C. et al., 219
Ganoza M.C. – See Kiel M.C. et al., 1097
Gärtner R. – See Bechtner G. et al., 315
Gaubert G., Gosselin G., Boudou V., Imbach J.L., Eriksson S., Maury G. – Low enantioselectivities of human deoxycytidine kinase and human deoxyguanosine kinase with respect to 2'-deoxyadenosine, 2'-deoxyguanosine and their analogs, 1041
Gaude T. – See Giranton J.L. et al., 675
Geel A.B.M. (van) – See Roos M. et al., 797
Gentil A. – See Le Page F. et al., 147
Gentile F. – See Veneziani B.M. et al., 517
Gerber H., Peter H.J., Bürgi U., Bigler S., Kaempf J., Zbaeren J. – Colloidal aggregates of insoluble inclusions in human goiters, 441
Giacomini P. – See Mischiati C. et al., 219
Giallauria F. – See Veneziani B.M. et al., 517
Giardina T., Biagini A., Massey-Harroche D., Puigserver A. – Distribution and subcellular localization of acylpeptide hydrolase and acylase I along the hog gastrointestinal tract, 1049
Giranton J.L., Passelègue E., Dumas C., Cock J.M., Gaude T. – Membrane proteins involved in pollen-pistil interactions, 675
Goffard J.C. – See Pirson I. et al., 309
Gómez I. – See Loyola C. et al., 261
Gómez M.J. – See Carrión M. et al., 879
Gomord V., Wee E., Faye L. – Protein retention and localization in the endoplasmic retikculum and the Golgi apparatus, 607
Gons M.H. – See Graaf S.A.R. (van de), 425
Gosselin G. – See Gaubert G. et al., 1041
Gouesbet G. – See Norris V. et al., 915
Graaf S.A.R. (van de), Cammenga M., Ponne

N.J., Veenboer G.J.M., Gons M.H., Orgiazzi J., Vijlder J.M.J. (de), Ris-Stalpers C. – The screening for mutations in the thyroglobulin cDNA from six patients with congenital hypothyroidism, 425
Grant N. – See Norris V. et al., 915
Gras R. – See Delamarche C. et al., 1065
Grassadonia A. – See Suzuki K. et al., 329
Grehan G. – See Norris V. et al., 915
Gressent F. – See Niebel A. et al., 669
Grienenberger J.M. – See Combettes B. and Grienenberger J.M., 645
Grignon C. – Recent advances on proteins of plant terminal membranes, 577
Grosjean H. – See Arluison V. et al., 751
Gross H.J. – See Fujiwara T. et al., 213
Guerdoux-Jamet P. – See Delamarche C. et al., 1065
Guespin J. – See Norris V. et al., 915
Guez V. – See Jermutus L. et al., 235
Günther Sillero M.A. – See Fontes R. et al., 229
Guzmán E.C. – See Molina F. et al., 811

H

Håkansson G. – See Escobar Galvis M.L. et al., 1089
Hanawalt P.C. – See Tornaletti S., Hanawalt P.C., 139
Handa N., Kobayashi I. – Post-segregational killing by restriction modification gene complexes: Observations of individual cell deaths, 931
Harney J.W. – See Salvatore D. et al., 535
Hartoft-Nielsen M.L. – See Rasmussen A.K. et al., 511
Hatipoglu B.A., Schneider A.B. – Selective endocytosis of thyroglobulin: A review of potential mechanisms for protecting newly synthesized molecules from premature degradation, 549
Hawes C. – See Satiat-Jeunemaitre B. et al., 597
Hemmersbach R. – See Meli A. et al., 281
Herrick J., Bensimon A. – Single molecule analysis of DNA replication, 859
Hilderman R.H., Fairbank A.T. – Binding and internalization of p^1,p^4-diadenosine 5'-tetraphosphate by bovine aortic endothelial cells, 255
Hjerpe A. – See Syrokou A. et al., 733
Hoeijmakers J.H.J. – See Boer J. de and Hoeijmakers J.H.J., 127
Holland I.B., Jones H.E., Campbell A.K., Jacq A. – An assessment of the role of intracellular free Ca^{2+} in *E. coli*, 901

I–J

Iantomasi T. – See Degl'Innocenti D. et al., 1025
Imbach J.L. – See Gaubert G. et al., 1041
Impiombato F.A. – See Meli A. et al., 281
Insinna E. – See Norris V. et al., 915
Iwai N. – See Wachi M. et al., 909
Jacq A. – See Holland I.B. et al., 901
Jermutus L., Guez V., Bedouelle H. – Disordered C-terminal domain of tyrosyl-tRNA synthetase: Secondary structure prediction, 235
Jiang P.H., Chany-Fournier F., Chany C. – Sarcolectin: Complete purification for molecular cloning, 701 – See Kaba A. et al., 709
Jiménez-Sánchez A. – See Molina F. et al., 811
Jones H.E. – See Holland I.B. et al., 901
Joyard J. – See Barbier-Brygoo H. and Joyard J., 575 – See Sato N. et al., 619

K

Kaba A., Jiang P.H., Chany-Fournier F., Chany C. – Sarcolectin (SCL): Structure and expression of the recombinant molecule, 709
Kaempf J. – See Gerber H. et al., 441
Karamanos N.K. – See Chatziioannidis C.C. et al., 187 – See Syrokou A. et al., 733
Karlsson M., Pavlov M.Y., Malmqvist M., Persson B., Ehrenberg M. – Initiation of *Escherichia coli* ribosomes on matrix coupled mRNAs studied by optical biosensor technique, 995
Katayama T., Sekimizu K. – Inactivation of *Escherichia coli* DnaA protein by DNA polymerase III and negative regulations for initiation of the chromosomal replication, 835
Kern R. – See d'Alençon E. et al., 841
Kieffer S. – See Tapparo A. et al., 943
Kiel M.C., Aoki H., Ganoza M.C. – Identification of a ribosomal ATPase in *Escherichia coli* cells, 1097
Kim W.B. – See Suzuki K. et al., 329
Kimura S., Ward J.M., Minoo P. – Thyroid-specific enhancer-binding protein/thyroid transcription factor 1 is not required for the initial specification of the thyroid and lung primordia, 321
Kjær S. – See Wind T. et al., 1079
Klaus C. – See Eckardt-Schupp F., Klaus C., 161
Klein G. – See Tapparo A. et al., 943
Klyman E. – See Binenbaum Z. et al., 921
Knorpp C. – See Escobar Galvis M.L. et al., 1089
Kobayashi I. – See Handa N. and Kobayashi I., 931
Kohiyama M. – See Landoulsi A. and Kohiyama M., 827 – See d'Alençon E. et al., 841
Kohn L.D. – See Suzuki K. et al., 329
Köhrle J. – The trace element selenium and the thyroid gland, 527
Kondo Y. – See Sho K. et al., 341
Koppes L.J.H., Woldringh C.L., Nanninga N. – *Escherichia coli* contains a DNA replication compartment in the cell center, 803
Kunihisa A. – See Wachi M. et al., 909
Kwon O.S., Churchich J.E. – Refolding of pyridoxine-5'-P oxidase assisted by GRoeL, 1057

L

Laezza C. – See Bifulco M. et al., 287
Landoulsi A., Kohiyama M. – Initiation of DNA replication in Δ*cya* mutants of *Escherichia coli* K12, 827
Lapi P. – See Macchia P.E. et al., 433
Larsen A.K. – See Scala D. et al., 771
Laugâa P. – See Bravo C. et al., 201
Lavaroni S. – See Suzuki K. et al., 329
Le Page F., Gentil A., Sarasin A. – Repair and mutagenesis survey of 8-hydroxyguanine in bacteria and human cells, 147
Le Sceller L. – See Norris V. et al., 915
Lebrun-Garcia A., Bourque S., Bineta M.N., Ouaked F., Wendehenne D., Chiltz A., Schäffner A., Pugin A. – Involvement of plasma membrane proteins in plant defense responses. Analysis of the cryptogein signal transduction in tobacco, 663
Lee J.C. – See Pakhomova O.N. et al., 1015
Legue O. – See Leseney A.M. et al., 373
Leroy A. – See Dassain M. et al., 889
Leseney A.M., Dème D., Legue O., Ohayon R., Chanson P., Sales J.P., Carvalho D.P., Dupuy C., Virion A. – Biochemical characterization of a Ca^{2+}/NAD(P)H-dependent H_2O_2 generator in human thyroid tissue, 373
Leuba S.H. – See An W. et al., 727
Li R.Y. – See Salles B. et al., 53
Liguoro D. – See Pacifico F. et al., 493
Lipardi C., Nitsch L., Zurzolo C. – Mechanisms of apical protein sorting in polarized thyroid epithelial cells, 347
Liu R.T. – See Takeda T. et al., 297
Loyola C., Saavedra C., Gómez I., Vásquez C. – The amino acidic substitution of cysteine 167 by serine ($C^{167}S$) in *Bst*VI restriction endonuclease of *Bacillus stearothermophilus* V affects its conformation and thermostability, 261
Lucisano A. – See Pelagalli A. et al., 1003

M

Macchia P.E., Mattei M.G., Lapi P., Fenzi G., Di Lauro R. – Cloning, chromosomal localization and identification of polymorphisms in the human thyroid transcription factor 2 gene (TITF2), 433
Maenhaut C. – See Pirson I. et al., 309
Maheu B. – See Norris V. et al., 915
Majka J. – See Messer W. et al., 819
Malmqvist M. – See Karlsson M. et al., 995
Mangani F. – See Piatti E. et al., 1011
Mansion M. – See Santonia V. et al., 655
Márquez J. – See Aledo J.C. et al., 1109
Martinis S.A., Plateau P., Cavarelli J., Florentz C. – Aminoacyl-tRNA synthetases: A new image for a classical family, 683
Marzano L.A. – See Vitale M. et al., 477
Marzocchini R. – See Degl'Innocenti D. et al., 1031
Masai H., Deneke J., Furui Y., Tanaka T., Arai K.I. – *Escherichia coli* and *Bacillus subtilis* PriA proteins essential for recombination-dependent DNA replication: Involvement of ATPase/helicase activity of PriA for inducible stable DNA replication, 847

Massey-Harroche D. – See Giardina T. et al., 1049
Mattei M.G. – See Macchia P.E. et al., 433
Maury G. – See Gaubert G. et al., 1041
Medeiros-Neto G. – See Tomimori E.K. et al., 447
Meli A., Perrella G., Curcio F., Hemmersbach R., Neubert J., Impiombato F.A. – Response to thyrotropin of normal thyroid follicular cell strain FRTL5 in hypergravity, 281
Melillo R.M. – See Santoro M. et al., 397
Ménissier-de Murcia J. – See Dantzer F. et al., 69
Merchante-Schubert R. – See Carrión M. et al., 879
Messer W., Blaesing F., Majka J., Nardmann J., Schaper S., Schmidt A., Seitz H., Speck C., Tüngler D., Wegrzyn G., Weigel C., Welzeck M., Zakrzewska-Czerwinska J. – Functional domains of DnaA proteins, 819
Minoo P. – See Kimura S. et al., 321
Mischiati C., Feriotto G., Bianchi N., Rutigliano C., Giacomini P., Gambari R. – Analysis of the human HLA-DRA gene upstream region: Evidence for a stem-loop array directed by nuclear factors, 219
Miyagi E. – See Suzuki K. et al., 329
Miyata M., Seto S. – Cell reproduction cycle of mycoplasma, 873
Mizutani T. – See Fujiwara T. et al., 213
Moggs J.G., Almouzni G. – Chromatin rearrangements during nucleotide excision repair, 45
Molina F., Jiménez-Sánchez A., Zyskind J.W., Guzmán E.C. – Chromosomal insertions localized around *oriC* affect the cell cycle in *Escherichia coli*, 811
Monette J. – See Pakhomova O.N. et al., 1015
Monnier C. – See Norris V. et al., 915
Montañez C. – See Bermúdez-Cruz R. et al., 757
Montuori N. – See Ragno P. et al., 355
Morassutti C., Scaggiante B., Dapas B., Xodo L., Tell G., Quadrifoglio F. – Effect of phosphorothioate modifications on the ability of GTn oligodeoxynucleotides to specifically recognize single-stranded DNA-binding proteins and to affect human cancer cellular growth, 1115
Mori A. – See Suzuki K. et al., 329
Morreale De Escobar G. – See Escobar-Morreale H.F. et al., 453
Mousli M. – See Brino L. et al., 973
Müeller F. – See Vitale M. et al., 477
Muller C., Calsou P., Frit P., Salles B. – Regulation of the DNA-dependent protein kinase (DNA-PK) activity in eucaryotic cells, 117

N

Nagai K. – See Wachi M. et al., 909
Nagasawa T. – See Takeda T. et al., 297
Nakazato M. – See Suzuki K. et al., 329
Nanninga N. – See Roos M. et al., 797 – See Koppes L.J.H. et al., 803
Nardmann J. – See Messer W. et al., 819
Narita T. – See Sho K. et al., 341
Neubert J. – See Meli A. et al., 281
Nicolas J. – See Delamarche C. et al., 1065
Niebel A., Gressent F., Bono J.J., Ranjeva R., Cullimore J. – Recent advances in the study of Nod factor perception and signal transduction, 669
Niedergang C. – See Dantzer F. et al., 69
Nitsch L. – See Lipardi C. et al., 347
Norris V., Alexandre S., Bouligand Y., Cellier D., Demarty M., Grehan G., Gouesbet G., Guespin J., Insinna E., Le Sceller L., Maheu B., Monnier C., Grant N., Onoda T., Orange N., Oshima A., Picton L., Polaert H., Ripoll C., Thellier M., Valleton J.M., Verdus M.C., Vincent J.C., White G., Wiggins P. – Hypothesis: Hyperstructures regulate bacterial structure and the cell cycle, 915

O

Obregón M.J. – See Escobar-Morreale H.F. et al., 453
Ohayon R. – See Leseney A.M. et al., 373
Ojha M., Cattaneo A., Schwendimann B. – Comparative studies of Ca^{2+}-dependent proteases (CDP I and CDP II) from *Allomyces arbuscula*, 765
Okajima F. – See Sho K. et al., 341
Oliver J. – See Dantzer F. et al., 69
Onoda T. – See Norris V. et al., 915
Oppenheimer J.H. – Evolving concepts of thyroid hormone action, 539
Orange N. – See Norris V. et al., 915
Orgiazzi J. – See Graaf S.A.R. (van de), 425
Oshima A. – See Norris V. et al., 915
Ouaked F. – See Lebrun-Garcia A. et al., 663
Oudet P. – See Brino L. et al., 973

P-Q

Pacifico F., Liguoro D., Acquaviva R., Formisano S., Consiglio E. – Thyroglobulin binding and TSH regulation of the RHL-1 subunit of the asialoglycoprotein receptor in rat thyroid, 493
Pacini F., Pinchera A. – Serum and tissue thyroglobulin measurement: Clinical applications in thyroid disease, 463
Pakhomova O.N., Yeh L.C.C., Monette J., Lee J.C. – In vitro assembly of yeast 5S rRNA and a fusion protein containing ribosomal protein L5 and maltose binding protein, 1015
Pandini G. – See Belfiore A. et al., 403
Pas E. – See Zaritsky A. et al., 897
Passelègue E. – See Giranton J.L. et al., 675
Pavlov M.Y. – See Karlsson M. et al., 995
Pelagalli A., Belisario M.A., Squillacioti C., Della Morte R., d'Angelo D., Tafuri S., Lucisano A., Staiano N. – The mycotoxin fumonisin B1 inhibits integrin-mediated cell-matrix adhesion, 1003
Perrella G. – See Meli A. et al., 281
Perrotte-Piquemal M., Danchin A., Biville F. – Pyrophosphate increases the efficiency of enterobactin-dependent iron uptake in *Escherichia coli*, 245
Persson B. – See Karlsson M. et al., 995
Peter H.J. – See Gerber H. et al., 441
Petit C., Sancar A. – Nucleotide excision repair: From *E. coli* to man, 15
Petric M. – See Eggo M.C. et al., 291
Piacentini M.P. – See Piatti E. et al., 1011
Piatti E., Piacentini M.P., Fraternale D., Bucchini A., Mangani F., Accorsi A. – *myo*-[^{3}H]-inositol loaded erythrocytes and white ghosts: Two models to investigate the phosphatidylinositol synthesis in human red cells, 1011
Picascia A.– See Chieffi P. et al., 367
Picton L. – See Norris V. et al., 915
Pietrarrelli M. – See Suzuki K. et al., 329
Pikula S. – See Danieluk M. et al., 717
Pinchera A. – See Pacini F. and Pinchera A., 463
Pirson I., Behrends J., Savonet V., Goffard J.C., Dumont J.E., Schurmans S., Maenhaut C. – Identification and characterization of mRNAs differentially expressed in thyroid cells stimulated by a mitogenic treatment, 309
Plateau P. – See Martinis S.A. et al., 683
Pohlenz J., Refetoff S. – Mutations in the sodium/iodide symporter (NIS) gene as a cause for iodide transport defects and congenital hypothyroidism, 469
Polaert H. – See Norris V. et al., 915
Polizio F. – See Santucci R. et al., 745
Ponne N.J. – See Graaf S.A.R. (van de), 425
Prost J.F., Cozzone A.J. – Detection of an extended –10 element in the promoter region of the *pckA* gene encoding phosphoenolpyruvate carboxykinase in *Escherichia coli*, 197
Pugin A. – See Lebrun-Garcia A. et al., 663
Puigserver A. – See Giardina T. et al., 1049
Quadrifoglio F. – See Morassutti C. et al., 1115

R

Rabilloud T. – See Santonia V. et al., 655
Radicella J.P. – See Boiteux S. and Radicella J.P., 59
Ragno P., Montuori N., Salzano S., Rossi G. – Urokinase-type plasminogen activator and normal thyroid cell adhesion to the extracellular matrix, 355
Ramponi G. – See Degl'Innocenti D. et al., 1025 – See Degl'Innocenti D. et al., 1031
Ranjeva R. – See Niebel A. et al., 669
Rasmussen A.K., Hartoft-Nielsen M.L., Feldt-Rasmussen U. – Models to study the pathogenesis of thyroid autoimmunity, 511
Raugei G. – See Degl'Innocenti D. et al., 1031
Reed Larsen P. – See Salvatore D. et al., 535
Refetoff S. – See Pohlenz J. and Refetoff S., 469
Riès-Kautt M. – See Arluison V. et al., 751
Ripoll C. – See Norris V. et al., 915
Ris-Stalpers C. – See Graaf S.A.R. (van de), 425
Robbins J. – See Benvenga S. and Robbins J., 545

Rodrigo G. – See Salles B. et al., 53
Rodríguez-Peña A. – See Escrivá H. et al., 965
Rolland N. – See Sato N. et al., 619
Rolli V. – See Dantzer F. et al., 69
Roos M., Geel A.B.M. (van), Aarsman M.E.G., Veuskens J.T.M., Woldringh C.L., Nanninga N. – Cellular localization of *oriC* during the cell cycle of *Escherichia coli* as analyzed by fluorescent in situ hybridization, 797
Ros P., Rossi D.L., Acebrón A., Santisteban P. – Thyroid-specific gene expression in the multi-step process of thyroid carcinogenesis, 389
Rosati F. – See Degl'Innocenti D. et al., 1025 – See Degl'Innocenti D. et al., 1031
Rossi D.L. – See Ros P. et al., 389
Rossi G. – See Ragno P. et al., 355 – See Vitale M. et al., 477
Rossignol M. – See Santonia V. et al., 655
Rouquié D. – See Santonia V. et al., 655
Ruijter G.J.G., Visser J. – Characterization of *Aspergillus niger* phosphoglucose isomerase. Use for quantitative determination of erythrose 4-phosphate, 267
Rutigliano C. – See Mischiati C. et al., 219

S

Saavedra C. – See Loyola C. et al., 261
Sachse A. – See Bechtner G. et al., 315
Saito J. – See Suzuki K. et al., 329
Sales J.P. – See Leseney A.M. et al., 373
Salles B., Rodrigo G., Li R.Y., Calsou P. – DNA damage excision repair in microplate wells with chemiluminescence detection: Development and perspectives, 53 – See Muller C. et al., 117
Salvatore D., Harney J.W., Reed Larsen P. – Mutation of the Secys residue 266 in human type 2 selenodeiodinase alters ^{75}Se incorporation without affecting its biochemical properties, 535
Salvatore G. – See Santoro M. et al., 397
Salvatore P. – See Chiariotti L. et al., 381
Salzano S. – See Ragno P. et al., 355 – See Vitale M. et al., 477
Sancar A. – See Petit C. and Sancar A., 15
Santisteban P. – See Ros P. et al., 389
Santonia V., Doumas P., Rouquié D., Mansion M., Rabilloud T., Rossignol M. – Large scale characterization of plant plasma membrane proteins, 655
Santoro M., Melillo R.M., Carlomagno F., Visconti R., De Vita G., Salvatore G., Fusco A., Vecchio G. – Different mutations of the RET gene cause different human tumoral diseases, 397
Santucci R., Polizio F., Desideri A. – Formation of a molten-globule-like state of cytochrome *c* induced by high concentrations of glycerol, 745
Sarasin A. – See Le Page F. et al., 147
Satiat-Jeunemaitre B., Boevink P., Hawes C. – Membrane trafficking in higher plant cells: GFP and antibodies, partners for probing the secretory pathway, 597
Sato N., Rolland N., Block M.A., Joyard J. – Do plastid envelope membranes play a role in the expression of the plastid genome?, 619
Satre M. – See Tapparo A. et al., 943
Savonet V. – See Pirson I. et al., 309
Scaggiante B. – See Morassutti C. et al., 1115
Scala D., Escargueil A.E., Couprie J., Larsen A.K. – The catalytic activities of DNA topoisomerase II are most closely associated with the DNA cleavage/religation steps, 771
Schäffner A. – See Lebrun-Garcia A. et al., 663
Schaper S. – See Messer W. et al., 819
Schild D. – See Thompson L.H. and Schild D., 87
Schmidt A. – See Messer W. et al., 819
Schneider A.B. – See Hatipoglu B.A. and Schneider A.B., 549
Schopohl D. – See Bechtner G. et al., 315
Schreiber V. – See Dantzer F. et al., 69
Schurmans S. – See Pirson I. et al., 309
Schwendimann B. – See Ojha M. et al., 765
Segura J.A. – See Aledo J.C. et al., 1109
Seitz H. – See Messer W. et al., 819
Sekimizu K. – See Katayama T. and Sekimizu K., 835
Seto S. – See Miyata M. and Seto S., 873
Shafran N. – See Suzuki K. et al., 329
Sho K., Narita T., Okajima F., Kondo Y. – An adenosine receptor agonist-induced modulation of TSH-dependent cell growth in FRTL-5 thyroid cells mediated by inhibitory G protein, G_i, 341
Sillero A. – See Fontes R. et al., 229
Speck C. – See Messer W. et al., 819
Squatrito S. – See Belfiore A. et al., 403
Squillacioti C. – See Pelagalli A. et al., 1003
Staiano N. – See Pelagalli A. et al., 1003
Stanzione R.– See Chieffi P. et al., 367
Stone S. – See Colzani R.M. et al., 485
Suzuki K., Mori A., Lavaroni S., Ulianich L., Miyagi E., Saito J., Nakazato M., Pietrarrelli M., Shafran N., Grassadonia A., Kim W.B., Consiglio E., Formisano S., Kohn L.D. – Thyroglobulin regulates follicular function and heterogeneity by suppressing thyroid-specific gene expression, 329
Suzuki S. – See Takeda T. et al., 297
Syrokou A., Tzanakakis G.N., Hjerpe A., Karamanos N.K. – Proteoglycans in human malignant mesothelioma. Stimulation of their synthesis induced by epidermal, insulin and platelet-derived growth factors involves receptors with tyrosine kinase activity, 733

T

Tafuri S. – See Pelagalli A. et al., 1003
Taghbalout A. – See d'Alençon E. et al., 841
Takeda T., Suzuki S., Nagasawa T., Liu R.T., DeGroot L.J. – DNA binding affinity of hTRb1 mutants as heterodimers with traps from different tissues, 297
Tanaka T. – See Masai H. et al., 847
Tarug P. – See Volpi N. and Tarug P., 955
Tata J.R. – Amphibian metamorphosis as a model for studying the developmental actions of thyroid hormone, 359
Taurog A. – Molecular evolution of thyroid peroxidase, 557
Tell G. – See Morassutti C. et al., 1115
Thacker J. – The role of homologous recombination processes in the repair of severe forms of DNA damage in mammalian cells, 77
The multigene immunophilin family of *Dictyostelium discoideum*. Tapparo A., Kieffer S., Cretin F., Satre M., Klein G. – Characterization of microsomal and mitochondrial cyclophilin isoforms, 943
Thellier M. – See Norris V. et al., 915
Thompson L.H., Schild D. – The contribution of homologous recombination in preserving genome integrity in mammalian cells, 87
Tomimori E.K., Camargo R.Y.A., Bisi H., Medeiros-Neto G. – Combined ultrasonographic and cytological studies in the diagnosis of thyroid nodules, 447
Tornaletti S., Hanawalt P.C. – Effect of DNA lesions on transcription elongation, 139
Tramontano D.– See Chieffi P. et al., 367
Trucco C. – See Dantzer F. et al., 69
Tsegenidis T. – See Chatziioannidis C.C. et al., 187
Tüngler D. – See Messer W. et al., 819
Tzanakakis G.N. – See Syrokou A. et al., 733

U–V

Ulianich L. – See Suzuki K. et al., 329
Vallejo C.G. – See Escrivá H. et al., 965
Valleton J.M. – See Norris V. et al., 915
Vallon O. – See de Vitry C. and Vallon O., 631
Van Den Hove M.F., Beckers C., Devlieger H., De Zegher F., De Nayer P. – Hormone synthesis and storage in the thyroid of human preterm and term newborns: effect of thyroxine treatment, 563
Van Geel A. – See Zaritsky A. et al., 897
van Holde K. – See An W. et al., 727
Vásquez C. – See Loyola C. et al., 261
Vecchio G. – See Santoro M. et al., 397
Veenboer G.J.M. – See Graaf S.A.R. (van de), 425
Vella V. – See Belfiore A. et al., 403
Veneziani B.M., Giallauria F., Gentile F. – The disulfide bond pattern between fragments obtained by the limited proteolysis of bovine thyroglobulin, 517
Verdus M.C. – See Norris V. et al., 915
Veuskens J.T.M. – See Roos M. et al., 797
Vigneri R. – See Belfiore A. et al., 403
Vijlder J.M.J. (de), – See Graaf S.A.R. (van de), 425
Villone G.– See Chieffi P. et al., 367
Vincent J.C. – See Norris V. et al., 915
Vincenzini M.T. – See Degl'Innocenti D. et al., 1025
Virion A. – See Leseney A.M. et al., 373
Visconti R. – See Santoro M. et al., 397
Visser J. – See Ruijter G.J.G. and Visser J., 267
Vitale M. , De Riu S., Fenzi G.F., Casamassima A., Salzano S., Müeller F., Marzano L.A.,

Rossi G. – Expression of integrins of the b family in thyroid cells from patients with Graves' disease in vivo and in vitro, 477
Vitale M.A. – See Eggo M.C. et al., 291
Volpi N., Tarug P. – The protective effect on Cu^{2+}- and AAPH-mediated oxidation of human low-density lipoproteins depends on glycosaminoglycan structure, 955
Vothknecht U.C. – See Das R. and Vothknecht U.C., 1037
Wachi M., Iwai N., Kunihisa A., Nagai K. – Irregular nuclear localization and anucleate cell production in *Escherichia coli* induced by a Ca^{2+} chelator, EGTA, 909

W

Ward J.M. – See Kimura S. et al., 321
Wee E. – See Gomord V. et al., 607
Wegrzyn G. – See Messer W. et al., 819
Weigel C. – See Messer W. et al., 819
Welzeck M. – See Messer W. et al., 819
Wendehenne D. – See Lebrun-Garcia A. et al., 663
White G. – See Norris V. et al., 915
Wiggins P. – See Norris V. et al., 915
Wind T., Kjær S., Clark B.F.C. – Display of Ras on filamentous phage through cysteine replacement, 1079
Woisard A. – See Bravo C. et al., 201
Woldringh C.L. – See Roos M. et al., 797 – See Koppes L.J.H. et al., 803 – See Zaritsky A. et al., 897
Wollman S.H. – Summary of observations on transplantable tumors of the rat thyroid gland, 409
Wood R.D. – DNA damage recognition during nucleotide excision repair in mammalian cells, 39

X–Z

Xodo L. – See Morassutti C. et al., 1115
Yeh L.C.C. – See Pakhomova O.N. et al., 1015
Zakrzewska-Czerwinska J. – See Messer W. et al., 819
Zaritsky A., Van Geel A., Fishov I., Pas E., Einav M., Woldringh C.L. – Visualizing multiple constrictions in spheroidal *Escherichia coli* cells, 897
Zbaeren J. – See Gerber H. et al., 441
Zdzienicka M.Z. – Mammalian X-ray-sensitive mutants which are defective in non-homologous (illegitimate) DNA double-strand break repair, 107
Zlatanova J. – See An W. et al., 727
Zurzolo C. – See Lipardi C. et al., 347
Zyskind J.W. – See Molina F. et al., 811

Biochimie 81 (1999) 1139–1148
© Société française de biochimie et biologie moléculaire / Elsevier, Paris

Keyword index 1999

A

acetylcholinesterase-similar domain – The disulfide bond pattern between fragments obtained by the limited proteolysis of bovine thyroglobulin, 517
actin – Membrane trafficking in higher plant cells: GFP and antibodies, partners for probing the secretory pathway, 597
acylase I – Distribution and subcellular localization of acylpeptide hydrolase and acylase I along the hog gastro-intestinal tract, 1049
acyl-CoA synthetase – Acyl-CoA synthetase catalyzes the synthesis of diadenosine hexaphosphate (Ap_6A), 229
acylpeptide hydrolase – Distribution and subcellular localization of acylpeptide hydrolase and acylase I along the hog gastro-intestinal tract, 1049
acylphosphatase – Acylphosphatase expression during macrophage differentiation and activation of U-937 cell line, 1031
adaptive response – Radiation inducible DNA repair processes in eukaryotes, 161
adenosine – An adenosine receptor agonist-induced modulation of TSH-dependent cell growth in FRTL-5 thyroid cells mediated by inhibitory G protein, G_i, 341
adhesion molecules – Expression of integrins of the β family in thyroid cells from patients with Graves' disease in vivo and in vitro, 477
aequorin – An assessment of the role of intracellular free Ca^{2+} in *E. coli*, 901
aggregation – RNA:pseudouridine synthetase Pus1 from *Saccharomyces cerevisiae*: Oligomerization property and stoichiometry of the complex with yeast $tRNA^{Phe}$, 751
aging – Cancer from the outside, aging from the inside: Mouse models to study the consequences of defective nucleotide excision repair, 127 – Oxidative DNA damage processing in nuclear and mitochondrial DNA, 155 – Aging, mitochondria, and coenzyme Q_{10}: The neglected relationship, 1131
Albert Kelner – The discovery of enzymatic photoreactivation and the question of priority: The letters of Salvador Luria and Albert Kelner, 7
Allomyces arbuscula – Comparative studies of Ca^{2+}-dependent proteases (CDP I and CDP II) from *Allomyces arbuscula*, 765
aminoacyl-tRNA synthetase – Disordered C-terminal domain of tyrosyl-tRNA synthetase: Secondary structure prediction, 235 – Aminoacyl-tRNA synthetases: A new image for a classical family, 683 – Phenylalanyl-tRNA synthetase from the archaeon *Methanobacterium thermoautotrophicum* is an $(\alpha\beta)_2$ heterotetrameric protein, 1037
animal models – Models to study the pathogenesis of thyroid autoimmunity, 511
animal peroxidases – Molecular evolution of thyroid peroxidase, 557
annexin VI – Annexin VI interacts with adenine nucleotides and their analogs, 717
antibiotics – Aminoacyl-tRNA synthetases: A new image for a classical family, 683 – Division-associated changes in membrane viscosity of *Escherichia coli*, 921
antioncogenes – Identification and characterization of mRNAs differentially expressed in thyroid cells stimulated by a mitogenic treatment, 309
anucleate cell – Irregular nuclear localization and anucleate cell production in *Escherichia coli* induced by a Ca^{2+} chelator, EGTA, 909
Ap_4A internalization – Binding and internalization of p^1,p^4-diadenosine 5'-tetraphosphate by bovine aortic endothelial cells, 255
Ap_4A receptor – Binding and internalization of p^1,p^4-diadenosine 5'-tetraphosphate by bovine aortic endothelial cells, 255
apical sorting – Mechanisms of apical protein sorting in polarized thyroid epithelial cells, 347
apoptosis – Transcription factor IIH: A key player in the cellular response to DNA damage, 27 – Induction of apoptosis in porcine thyroid follicles by transforming growth factor β1 and epidermal growth factor, 315
Arabidopsis thaliana – Large scale characterization of plant plasma membrane proteins, 655
arachidonic acid – Signaling pathways involved in thyroid hyperfunction and growth in Graves' disease, 415
archaea – Phenylalanyl-tRNA synthetase from the archaeon *Methanobacterium thermoautotrophicum* is an $(\alpha\beta)_2$ heterotetrameric protein, 1037
Aspergillus niger – Characterization of *Aspergillus niger* phosphoglucose isomerase. Use for quantitative determination of erythrose 4-phosphate, 267
A-state – Formation of a molten-globule-like state of cytochrome *c* induced by high concentrations of glycerol, 745
ATP – Annexin VI interacts with adenine nucleotides and their analogs, 717
ATP binding – Isoleucine 10 is essential for DNA gyrase B function in *Escherichia coli*, 973
ATP hydrolysis – The catalytic activities of DNA topoisomerase II are most closely associated with the DNA cleavage/religation steps, 771
ATPase – Isoleucine 10 is essential for DNA gyrase B function in *Escherichia coli*, 973 – Identification of a ribosomal ATPase in *Escherichia coli* cells, 1097
attachment organelle – Cell reproduction cycle of mycoplasma, 873
autoimmune thyroid disease – Models to study the pathogenesis of thyroid autoimmunity, 511
autoinduction – Amphibian metamorphosis as a model for studying the developmental actions of thyroid hormone, 359
autoradiography – *Escherichia coli* contains a DNA replication compartment in the cell center, 803

B

B. subtilis – An assessment of the role of intracellular free Ca^{2+} in *E. coli*, 901
bacterial membrane – Division-associated changes in membrane viscosity of *Escherichia coli*, 921
bacterial promoter – Detection of an extended –10 element in the promoter region of the *pckA* gene encoding phosphoenolpyruvate carboxykinase in *Escherichia coli*, 197
base excision repair – Involvement of poly(ADP-ribose) polymerase in base excision repair, 69 – Oxidative DNA damage processing in nuclear and mitochondrial DNA, 155
Becker muscular dystrophy – GSH system in relation to redox state in dystrophic skin fibroblasts, 1025
bending – Chromosomal insertions localized around *oriC* affect the cell cycle in *Escherichia coli*, 811
BIACORE – Initiation of *Escherichia coli* ribosomes on matrix coupled mRNAs studied by optical biosensor technique, 995
biochemical properties – Mutation of the Secys residue 266 in human type 2 selenodeiodinase alters ^{75}Se incorporation without affecting its biochemical properties, 535
bioenergetics – Aging, mitochondria, and coenzyme Q_{10}: The neglected relationship, 1131
blood capillaries – Summary of observations on transplantable tumors of the rat thyroid gland, 409
breast cancer – Insulin/IGF-I hybrid receptors

play a major role in IGF-I signaling in thyroid cancer, 403
brefeldin A – Membrane trafficking in higher plant cells: GFP and antibodies, partners for probing the secretory pathway, 597

C

Ca^{2+} – Irregular nuclear localization and anucleate cell production in *Escherichia coli* induced by a Ca^{2+} chelator, EGTA, 909
Ca^{2+}-channels – An assessment of the role of intracellular free Ca^{2+} in *E. coli*, 901
calcium – Annexin VI interacts with adenine nucleotides and their analogs, 717 – An assessment of the role of intracellular free Ca^{2+} in *E. coli*, 901
cAMP – Response to thyrotropin of normal thyroid follicular cell strain FRTL5 in hypergravity, 281
cancer – Identification and characterization of mRNAs differentially expressed in thyroid cells stimulated by a mitogenic treatment, 309
cancer therapy – Regulation of the DNA-dependent protein kinase (DNA-PK) activity in eucaryotic cells, 117
carbohydrate – Depletion of divalent cations within the secretory pathway inhibits the terminal glycosylation of complex carbohydrates of thyroglobulin, 497
carcinoma – Different mutations of the RET gene cause different human tumoral diseases, 397
catalytic activity – The catalytic activities of DNA topoisomerase II are most closely associated with the DNA cleavage/religation steps, 771
cDNA – Cloning and characterisation of a pea mitochondrial NDPK, 1089
CDP I and CDP II – Comparative studies of Ca^{2+}-dependent proteases (CDP I and CDP II) from *Allomyces arbuscula*, 765
cell adhesion – Urokinase-type plasminogen activator and normal thyroid cell adhesion to the extracellular matrix, 355 – The mycotoxin fumonisin B1 inhibits integrin-mediated cell-matrix adhesion, 1003
cell cycle – Transcription factor IIH: A key player in the cellular response to DNA damage, 27 – An adenosine receptor agonist-induced modulation of TSH-dependent cell growth in FRTL-5 thyroid cells mediated by inhibitory G protein, G_i, 341 – Inactivation of *Escherichia coli* DnaA protein by DNA polymerase III and negative regulations for initiation of the chromosomal replication, 835 – An assessment of the role of intracellular free Ca^{2+} in *E. coli*, 901 – Hypothesis: Hyperstructures regulate bacterial structure and the cell cycle, 915
cell death – Post-segregational killing by restriction modification gene complexes: Observations of individual cell deaths, 931
cell defence – The trace element selenium and the thyroid gland, 527
cell division – Cell reproduction cycle of mycoplasma, 873 – *mraW*, an essential gene at the *dcw* cluster of *Escherichia coli* codes for a cytoplasmic protein with methyltransferase activity, 879 – A new essential gene of the 'minimal genome' affecting cell division, 889 – An assessment of the role of intracellular free Ca^{2+} in *E. coli*, 901 – Division-associated changes in membrane viscosity of *Escherichia coli*, 921
cell extracts – DNA damage excision repair in microplate wells with chemiluminescence detection: Development and perspectives, 53
cell growth – Endogenous insulin-like growth factors regulate the proliferation of TSH-independent mutants derived from FRTL5 cells, 367 – The mycotoxin fumonisin B1 inhibits integrin-mediated cell-matrix adhesion, 1003
checkpoints – DNA structure checkpoint pathways in *Schizosaccharomyces pombe*, 173 – A new essential gene of the 'minimal genome' affecting cell division, 889
chemical cross-linking – Annexin VI interacts with adenine nucleotides and their analogs, 717
chemiluminescence – DNA damage excision repair in microplate wells with chemiluminescence detection: Development and perspectives, 53
Chlamydomonas – Mutants of *Chlamydomonas*: Tools to study thylakoid membrane structure, function and biogenesis, 631
chloroplast DNA – Do plastid envelope membranes play a role in the expression of the plastid genome?, 619
chondroitin – The protective effect on Cu^{2+}- and AAPH-mediated oxidation of human low-density lipoproteins depends on glycosaminoglycan structure, 955
chromatin rearrangements – Chromatin rearrangements during nucleotide excision repair, 45
chromosome partitioning – Irregular nuclear localization and anucleate cell production in *Escherichia coli* induced by a Ca^{2+} chelator, EGTA, 909
Cibacron blue 3GA – Annexin VI interacts with adenine nucleotides and their analogs, 717
circular dichroism – Formation of a molten-globule-like state of cytochrome *c* induced by high concentrations of glycerol, 745 – Refolding of pyridoxine-5'-P oxidase assisted by GRoeL, 1057
cleavage activity – A Y form of hammerhead ribozyme trapped by photo-cross-links retains full cleavage activity, 201
Cockayne syndrome – Oxidative DNA damage processing in nuclear and mitochondrial DNA, 155
coenzyme Q_{10} – Aging, mitochondria, and coenzyme Q_{10}: The neglected relationship, 1131
colloid – Colloidal aggregates of insoluble inclusions in human goiters, 441
complex I – Analysis of wheat mitochondrial complex I purified by a one-step immunoaffinity chromatography, 645
computational analysis – A symbolic-numeric approach to find patterns in genomes. Application to the translation initiation sites of *E. coli*, 1065
congenital hypothyroidism – The screening for mutations in the thyroglobulin cDNA from six patients with congenital hypothyroidism, 425 – Cloning, chromosomal localization and identification of polymorphisms in the human thyroid transcription factor 2 gene (TITF2), 433 – Mutations in the sodium/iodide symporter (NIS) gene as a cause for iodide transport defects and congenital hypothyroidism, 469
connective tissue – Summary of observations on transplantable tumors of the rat thyroid gland, 409
cooperativity – Functional domains of DnaA proteins, 819
core histone tails – The site of binding of linker histone to the nucleosome does not depend upon the amino termini of core histones, 727
covalent DNA-protein complexes – The catalytic activities of DNA topoisomerase II are most closely associated with the DNA cleavage/religation steps, 771
cyclic AMP – Initiation of DNA replication in Δ*cya* mutants of *Escherichia coli* K12, 827
cyclophilin – Characterization of microsomal and mitochondrial cyclophilin isoforms, 943
cysteine-rich repeats – The disulfide bond pattern between fragments obtained by the limited proteolysis of bovine thyroglobulin, 517
cytochrome $b_6 f$ – Mutants of *Chlamydomonas*: Tools to study thylakoid membrane structure, function and biogenesis, 631
cytochrome *c* – Formation of a molten-globule-like state of cytochrome *c* induced by high concentrations of glycerol, 745
cytokines – Models to study the pathogenesis of thyroid autoimmunity, 511
cytoskeleton – Cell reproduction cycle of mycoplasma, 873
cytotoxicity – Effect of phosphorothioate modifications on the ability of GTn oligodeoxynucleotides to specifically recognize single-stranded DNA-binding proteins and to affect human cancer cellular growth, 1115

D

damage recognition – DNA damage recognition during nucleotide excision repair in mammalian cells, 39
***dcw* cluster** – *mraW*, an essential gene at the *dcw* cluster of *Escherichia coli* codes for a cytoplasmic protein with methyltransferase activity, 879

Δ*cya* mutant – Initiation of DNA replication in Δ*cya* mutants of *Escherichia coli* K12, 827
deiodinase – The trace element selenium and the thyroid gland, 527
deoxycytidine kinase – Low enantioselectivities of human deoxycytidine kinase and human deoxyguanosine kinase with respect to 2'-deoxyadenosine, 2'-deoxyguanosine and their analogs, 1041
deoxyguanosine kinase – Low enantioselectivities of human deoxycytidine kinase and human deoxyguanosine kinase with respect to 2'-deoxyadenosine, 2'-deoxyguanosine and their analogs, 1041
dermatan – The protective effect on Cu^{2+}- and AAPH-mediated oxidation of human low-density lipoproteins depends on glycosaminoglycan structure, 955
dermatan sulphate – Purification and characterisation of a minor low-sulphated dermatan sulphate-proteoglycan from ray skin, 187
development – Characterization of microsomal and mitochondrial cyclophilin isoforms, 943
diadenosine hexaphosphate – Acyl-CoA synthetase catalyzes the synthesis of diadenosine hexaphosphate (Ap_6A), 229
diadenosine pentaphosphate – Acyl-CoA synthetase catalyzes the synthesis of diadenosine hexaphosphate (Ap_6A), 229
Dictyostelium discoideum – Characterization of microsomal and mitochondrial cyclophilin isoforms, 943
differentiation – Thyroid-specific gene expression in the multi-step process of thyroid carcinogenesis, 389 – Acylphosphatase expression during macrophage differentiation and activation of U-937 cell line, 1031
dinucleoside polyphosphates – Acyl-CoA synthetase catalyzes the synthesis of diadenosine hexaphosphate (Ap_6A), 229
disulfide bond – The disulfide bond pattern between fragments obtained by the limited proteolysis of bovine thyroglobulin, 517
disulphide bridge – Display of Ras on filamentous phage through cysteine replacement, 1079
divalent cations – Depletion of divalent cations within the secretory pathway inhibits the terminal glycosylation of complex carbohydrates of thyroglobulin, 497
division planes – Visualizing multiple constrictions in spheroidal *Escherichia coli* cells, 897
D-loop – *Escherichia coli* and *Bacillus subtilis* PriA proteins essential for recombination-dependent DNA replication: Involvement of ATPase/helicase activity of PriA for inducible stable DNA replication, 847
DNA binding affinity – DNA binding affinity of hTRβ1 mutants as heterodimers with traps from different tissues, 297
DNA binding protein – Involvement of poly(ADP-ribose) polymerase in base excision repair, 69
DNA damage – Nucleotide excision repair: From *E. coli* to man, 15 – DNA damage excision repair in microplate wells with chemiluminescence detection: Development and perspectives, 53 – DNA structure checkpoint pathways in *Schizosaccharomyces pombe,* 173
DNA helicase – *Escherichia coli* and *Bacillus subtilis* PriA proteins essential for recombination-dependent DNA replication: Involvement of ATPase/helicase activity of PriA for inducible stable DNA replication, 847
DNA imaging – Single molecule analysis of DNA replication, 859
DNA methylation – Post-segregational killing by restriction modification gene complexes: Observations of individual cell deaths, 931
DNA polymerase – Inactivation of *Escherichia coli* DnaA protein by DNA polymerase III and negative regulations for initiation of the chromosomal replication, 835
DNA recombination – *Escherichia coli* and *Bacillus subtilis* PriA proteins essential for recombination-dependent DNA replication: Involvement of ATPase/helicase activity of PriA for inducible stable DNA replication, 847
DNA repair – The discovery of enzymatic photoreactivation and the question of priority: The letters of Salvador Luria and Albert Kelner, 7
DNA repair – Nucleotide excision repair: From *E. coli* to man, 15 – Transcription factor IIH: A key player in the cellular response to DNA damage, 27 – DNA damage recognition during nucleotide excision repair in mammalian cells, 39 – Base excision repair of 8-hydroxyguanine protects DNA from endogenous oxidative stress, 59 – The role of homologous recombination processes in the repair of severe forms of DNA damage in mammalian cells, 77 – Repair and mutagenesis survey of 8-hydroxyguanine in bacteria and human cells, 147 – Oxidative DNA damage processing in nuclear and mitochondrial DNA, 155
DNA replication – DNA structure checkpoint pathways in *Schizosaccharomyces pombe,* 173 – *Escherichia coli* contains a DNA replication compartment in the cell center, 803 – Inactivation of *Escherichia coli* DnaA protein by DNA polymerase III and negative regulations for initiation of the chromosomal replication, 835 – *Escherichia coli* and *Bacillus subtilis* PriA proteins essential for recombination-dependent DNA replication: Involvement of ATPase/helicase activity of PriA for inducible stable DNA replication, 847 – Single molecule analysis of DNA replication, 859 – Cell reproduction cycle of mycoplasma, 873
DnaA – Inactivation of *Escherichia coli* DnaA protein by DNA polymerase III and negative regulations for initiation of the chromosomal replication, 835
DnaA protein – Initiation of DNA replication in Δ*cya* mutants of *Escherichia coli* K12, 827
DnaB – Functional domains of DnaA proteins, 819
DNA-binding protein – Do plastid envelope membranes play a role in the expression of the plastid genome?, 619
DNA-gyrase – Isoleucine 10 is essential for DNA gyrase B function in *Escherichia coli*, 973
DNA-PK – Regulation of the DNA-dependent protein kinase (DNA-PK) activity in eucaryotic cells, 117
dodecyl-—D-maltoside – RNA:pseudouridine synthetase Pus1 from *Saccharomyces cerevisiae*: Oligomerization property and stoichiometry of the complex with yeast $tRNA^{Phe}$, 751
dominant negative effect – DNA binding affinity of hTRβ1 mutants as heterodimers with traps from different tissues, 297
double-strand breaks – The contribution of homologous recombination in preserving genome integrity in mammalian cells, 87
DSB repair – Mammalian X-ray-sensitive mutants which are defective in non-homologous (illegitimate) DNA double-strand break repair, 107
Duchenne muscular dystrophy – GSH system in relation to redox state in dystrophic skin fibroblasts, 1025

E

EF-3 – Identification of a ribosomal ATPase in *Escherichia coli* cells, 1097
EGTA – Irregular nuclear localization and anucleate cell production in *Escherichia coli* induced by a Ca^{2+} chelator, EGTA, 909
Ehrlich ascites cell – Upregulation of glyceraldehyde-3-phosphate dehydrogenase mRNA in the spleen of tumor-bearing mice, 1109
elasticity – The structures of elastins and their function, 981
elastin – The structures of elastins and their function, 981
elicitor – Involvement of plasma membrane proteins in plant defense responses. Analysis of the cryptogein signal transduction in tobacco, 663
enantioselectivity – Low enantioselectivities of human deoxycytidine kinase and human deoxyguanosine kinase with respect to 2'-deoxyadenosine, 2'-deoxyguanosine and their analogs, 1041
endocytosis – Thyroglobulin binding and TSH regulation of the RHL-1 subunit of the asialoglycoprotein receptor in rat thy-

roid, 493 – Selective endocytosis of thyroglobulin: A review of potential mechanisms for protecting newly synthesized molecules from premature degradation, 549

endogenous oxidative stress – Base excision repair of 8-hydroxyguanine protects DNA from endogenous oxidative stress, 59

endoplasmic reticulum – Protein retention and localization in the endoplasmic reticulum and the Golgi apparatus, 607

endothelial cells – Binding and internalization of p^1,p^4-diadenosine 5'-tetraphosphate by bovine aortic endothelial cells, 255

enzymatic systems – Recent advances on proteins of plant terminal membranes, 577

epidermal growth factor – Induction of apoptosis in porcine thyroid follicles by transforming growth factor β1 and epidermal growth factor, 315

epigenetics – Post-segregational killing by restriction modification gene complexes: Observations of individual cell deaths, 931

erythrocyte – *myo*-[^{3}H]-inositol loaded erythrocytes and white ghosts: Two models to investigate the phosphatidylinositol synthesis in human red cells, 1011

erythrose 4-phosphate – Characterization of *Aspergillus niger* phosphoglucose isomerase. Use for quantitative determination of erythrose 4-phosphate, 267

Escherichia coli – Cellular localization of *oriC* during the cell cycle of *Escherichia coli* as analyzed by fluorescent in situ hybridization, 797 – *Escherichia coli* contains a DNA replication compartment in the cell center, 803 – Chromosomal insertions localized around *oriC* affect the cell cycle in *Escherichia coli*, 811 – *mraW*, an essential gene at the *dcw* cluster of *Escherichia coli* codes for a cytoplasmic protein with methyltransferase activity, 879 – An assessment of the role of intracellular free Ca^{2+} in *E. coli*, 901 – Irregular nuclear localization and anucleate cell production in *Escherichia coli* induced by a Ca^{2+} chelator, EGTA, 909 – Isoleucine 10 is essential for DNA gyrase B function in *Escherichia coli*, 973

evolution – Molecular evolution of thyroid peroxidase, 557

excision nuclease – Nucleotide excision repair: From *E. coli* to man, 15

excision repair – DNA damage excision repair in microplate wells with chemiluminescence detection: Development and perspectives, 53

extended –10 element – Detection of an extended –10 element in the promoter region of the *pckA* gene encoding phosphoenolpyruvate carboxykinase in *Escherichia coli*, 197

F

farnesylation inhibition – Inhibition of farnesylation blocks growth but not differentiation in FRTL-5 thyroid cells, 287

fetus – Effects of iodine repletion on thyroid morphology in iodine and/or selenium deficient rat term fetuses, pups and mothers, 485

fine needle biopsy – Combined ultrasonographic and cytological studies in the diagnosis of thyroid nodules, 447

Fischer 344 rat – Summary of observations on transplantable tumors of the rat thyroid gland, 409

fluorescence – Refolding of pyridoxine-5'-P oxidase assisted by GRoeL, 1057

fluorescence anisotropy – Division-associated changes in membrane viscosity of *Escherichia coli*, 921

fluorescent in situ hybridization – Cellular localization of *oriC* during the cell cycle of *Escherichia coli* as analyzed by fluorescent in situ hybridization, 797

fluorescent microscopy – Visualizing multiple constrictions in spheroidal *Escherichia coli* cells, 897

FMN – Refolding of pyridoxine-5'-P oxidase assisted by GRoeL, 1057

follicular function – Thyroglobulin regulates follicular function and heterogeneity by suppressing thyroid-specific gene expression, 329

FRT cell line – Mechanisms of apical protein sorting in polarized thyroid epithelial cells, 347

FRTL5 cells – Endogenous insulin-like growth factors regulate the proliferation of TSH-independent mutants derived from FRTL5 cells, 367

FtsZ – A new essential gene of the 'minimal genome' affecting cell division, 889 – An assessment of the role of intracellular free Ca^{2+} in *E. coli*, 901

FtsZ rings and arcs – Visualizing multiple constrictions in spheroidal *Escherichia coli* cells, 897

fumonisin B1 – The mycotoxin fumonisin B1 inhibits integrin-mediated cell-matrix adhesion, 1003

fusion protein – In vitro assembly of yeast 5S rRNA and a fusion protein containing ribosomal protein L5 and maltose binding protein, 1015

G

galectins – Control of galectin gene expression, 381

GAPDH – Upregulation of glyceraldehyde-3-phosphate dehydrogenase mRNA in the spleen of tumor-bearing mice, 1109

gene expression – Control of galectin gene expression, 381

genetic instability – Cancer from the outside, aging from the inside: Mouse models to study the consequences of defective nucleotide excision repair, 127

genetic stability – The contribution of homologous recombination in preserving genome integrity in mammalian cells, 87

genetics – The screening for mutations in the thyroglobulin cDNA from six patients with congenital hypothyroidism, 425

genome of *E. coli* – A symbolic-numeric approach to find patterns in genomes. Application to the translation initiation sites of *E. coli*, 1065

genome sequences – Phenylalanyl-tRNA synthetase from the archaeon *Methanobacterium thermoautotrophicum* is an $(\alpha\beta)_2$ heterotetrameric protein, 1037

genomics – A new essential gene of the 'minimal genome' affecting cell division, 889

GFP – Membrane trafficking in higher plant cells: GFP and antibodies, partners for probing the secretory pathway, 597

glutamate dehydrogenase – Identification of lysine residue involved in inactivation of brain glutamate dehydrogenase isoproteins by o-phthalaldehyde, 1123

glutathione – GSH system in relation to redox state in dystrophic skin fibroblasts, 1025

glycerol – Formation of a molten-globule-like state of cytochrome *c* induced by high concentrations of glycerol, 745

glycosaminoglycans – Proteoglycans in human malignant mesothelioma. Stimulation of their synthesis induced by epidermal, insulin and platelet-derived growth factors involves receptors with tyrosine kinase activity, 733

goiter – Colloidal aggregates of insoluble inclusions in human goiters, 441

Golgi apparatus – Protein retention and localization in the endoplasmic reticulum and the Golgi apparatus, 607

Graves' disease – Signaling pathways involved in thyroid hyperfunction and growth in Graves' disease, 415 – Expression of integrins of the β family in thyroid cells from patients with Graves' disease in vivo and in vitro, 477

GroEL – Refolding of pyridoxine-5'-P oxidase assisted by GRoeL, 1057

growth factors – Proteoglycans in human malignant mesothelioma. Stimulation of their synthesis induced by epidermal, insulin and platelet-derived growth factors involves receptors with tyrosine kinase activity, 733

GSH-related enzymes – GSH system in relation to redox state in dystrophic skin fibroblasts, 1025

GTPase – A new essential gene of the 'minimal genome' affecting cell division, 889

H

8-hydroxyguanine – Base excision repair of 8-hydroxyguanine protects DNA from endogenous oxidative stress, 59 – Repair and mutagenesis survey of 8-hydroxyguanine in bacteria and human cells, 147

H_2O_2 – Biochemical characterization of a Ca^{2+}/NAD(P)H-dependent H_2O_2 generator in human thyroid tissue, 373

H3 tails – The site of binding of linker histone

to the nucleosome does not depend upon the amino termini of core histones, 727
hammerhead ribozyme – A Y form of hammerhead ribozyme trapped by photocross-links retains full cleavage activity, 201
heparan – The protective effect on Cu^{2+}- and AAPH-mediated oxidation of human low-density lipoproteins depends on glycosaminoglycan structure, 955
heparin – The protective effect on Cu^{2+}- and AAPH-mediated oxidation of human low-density lipoproteins depends on glycosaminoglycan structure, 955
histology – Summary of observations on transplantable tumors of the rat thyroid gland, 409
history of science – The discovery of enzymatic photoreactivation and the question of priority: The letters of Salvador Luria and Albert Kelner, 7
HLA-DRA gene – Analysis of the human HLA-DRA gene upstream region: Evidence for a stem-loop array directed by nuclear factors, 219
HMG-CoA – Inhibition of farnesylation blocks growth but not differentiation in FRTL-5 thyroid cells, 287
hog enterocyte – Distribution and subcellular localization of acylpeptide hydrolase and acylase I along the hog gastro-intestinal tract, 1049
homeostasis – Tissue-specific patterns of changes in 3,5,3´-triiodo-L-thyronine concentrations in thyroidectomized rats infused with increasing doses of the hormone. Which are the regulatory mechanisms?, 453
homologous recombination – The role of homologous recombination processes in the repair of severe forms of DNA damage in mammalian cells, 77
human repair syndromes – Cancer from the outside, aging from the inside: Mouse models to study the consequences of defective nucleotide excision repair, 127
human T-lymphoblasts – Effect of phosphorothioate modifications on the ability of GTn oligodeoxynucleotides to specifically recognize single-stranded DNA-binding proteins and to affect human cancer cellular growth, 1115
hydrogen peroxide – Biochemical characterization of a Ca^{2+}/NAD(P)H-dependent H_2O_2 generator in human thyroid tissue, 373
hypergravity – Response to thyrotropin of normal thyroid follicular cell strain FRTL5 in hypergravity, 281

I

IGF-I receptor – Insulin/IGF-I hybrid receptors play a major role in IGF-I signaling in thyroid cancer, 403
immune response – Upregulation of glyceraldehyde-3-phosphate dehydrogenase mRNA in the spleen of tumor-bearing mice, 1109
immune system – Aging, mitochondria, and coenzyme Q_{10}: The neglected relationship, 1131
immunoaffinity – Analysis of wheat mitochondrial complex I purified by a one-step immunoaffinity chromatography, 645
immunocytochemistry – Membrane trafficking in higher plant cells: GFP and antibodies, partners for probing the secretory pathway, 597
immunophilin – Characterization of microsomal and mitochondrial cyclophilin isoforms, 943
in vitro models – Models to study the pathogenesis of thyroid autoimmunity, 511
in vitro protein synthesis – Initiation of *Escherichia coli* ribosomes on matrix coupled mRNAs studied by optical biosensor technique, 995
inactivation – Comparative studies of Ca^{2+}-dependent proteases (CDP I and CDP II) from *Allomyces arbuscula*, 765
inducible DNA-repair – Radiation inducible DNA repair processes in eukaryotes, 161
infectivity – Effects of thyroid-stimulating hormone on the replicative cycle of vesicular stomatitis virus in primary cultured sheep thyroid cells, 291
initiation – Initiation of DNA replication in Δ*cya* mutants of *Escherichia coli* K12, 827 – Initiation of *Escherichia coli* ribosomes on matrix coupled mRNAs studied by optical biosensor technique, 995
initiation of transcription – Detection of an extended –10 element in the promoter region of the *pckA* gene encoding phosphoenolpyruvate carboxykinase in *Escherichia coli*, 197
insulin receptor – Insulin/IGF-I hybrid receptors play a major role in IGF-I signaling in thyroid cancer, 403
insulin/IGF-I hybrid receptors – Insulin/IGF-I hybrid receptors play a major role in IGF-I signaling in thyroid cancer, 403
insulin-like growth factors – Insulin/IGF-I hybrid receptors play a major role in IGF-I signaling in thyroid cancer, 403
integrin receptors – The mycotoxin fumonisin B1 inhibits integrin-mediated cell-matrix adhesion, 1003
integrins – Expression of integrins of the β family in thyroid cells from patients with Graves' disease in vivo and in vitro, 477
interstrand cross-links – The contribution of homologous recombination in preserving genome integrity in mammalian cells, 87
intrinsic fluorescence – Annexin VI interacts with adenine nucleotides and their analogs, 717
iodide transport – Mutations in the sodium/iodide symporter (NIS) gene as a cause for iodide transport defects and congenital hypothyroidism, 469
iodine – Colloidal aggregates of insoluble inclusions in human goiters, 441 – Effects of iodine repletion on thyroid morphology in iodine and/or selenium deficient rat term fetuses, pups and mothers, 485 – The importance of thyroglobulin structure for thyroid hormone biosynthesis, 505 – The trace element selenium and the thyroid gland, 527
iodothyronine deiodinases – Tissue-specific patterns of changes in 3,5,3´-triiodo-L-thyronine concentrations in thyroidectomized rats infused with increasing doses of the hormone. Which are the regulatory mechanisms?, 453

K–L

kinase – Different mutations of the RET gene cause different human tumoral diseases, 397
kinetics – Characterization of *Aspergillus niger* phosphoglucose isomerase. Use for quantitative determination of erythrose 4-phosphate, 267
λ *tI* – Nus A is involved in transcriptional termination on λ *tI*, 757
limited proteolysis – The disulfide bond pattern between fragments obtained by the limited proteolysis of bovine thyroglobulin, 517
linker histone – The site of binding of linker histone to the nucleosome does not depend upon the amino termini of core histones, 727
lipid raft – Mechanisms of apical protein sorting in polarized thyroid epithelial cells, **347**
lipochitooligosaccharides – Recent advances in the study of Nod factor perception and signal transduction, 669
liquid crystal – Hypothesis: Hyperstructures regulate bacterial structure and the cell cycle, 915
L-nucleoside analogs – Low enantioselectivities of human deoxycytidine kinase and human deoxyguanosine kinase with respect to 2'-deoxyadenosine, 2'-deoxyguanosine and their analogs, 1041
loading – *myo*-[^{3}H]-inositol loaded erythrocytes and white ghosts: Two models to investigate the phosphatidylinositol synthesis in human red cells, 1011
low radiation doses – Radiation inducible DNA repair processes in eukaryotes, 161
low-density lipoprotein – The protective effect on Cu^{2+}- and AAPH-mediated oxidation of human low-density lipoproteins depends on glycosaminoglycan structure, 955
lung morphogenesis – Thyroid-specific enhancer-binding protein/thyroid transcription factor 1 is not required for the initial specification of the thyroid and lung primordia, 321

M

mammalian cells – The role of homologous recombination processes in the repair of

severe forms of DNA damage in mammalian cells, 77 – Radiation inducible DNA repair processes in eukaryotes, 161
mecillinam – Visualizing multiple constrictions in spheroidal *Escherichia coli* cells, 897
membrane – *myo*-[^{3}H]-inositol loaded erythrocytes and white ghosts: Two models to investigate the phosphatidylinositol synthesis in human red cells, 1011
membrane domains – Division-associated changes in membrane viscosity of *Escherichia coli*, 921
membrane *oriC* sequestration – Replication cycle dependent association of SeqA to the outer membrane fraction of *E. coli*, 841
membrane proteins – Protein retention and localization in the endoplasmic reticulum and the Golgi apparatus, 607 – Mutants of *Chlamydomonas*: Tools to study thylakoid membrane structure, function and biogenesis, 631
membrane viscosity – Division-associated changes in membrane viscosity of *Escherichia coli*, 921
mesothelioma – Proteoglycans in human malignant mesothelioma. Stimulation of their synthesis induced by epidermal, insulin and platelet-derived growth factors involves receptors with tyrosine kinase activity, 733
messenger and protein steady-state levels – Expression of mitochondrial genes and of the transcription factors involved in the biogenesis of mitochondria Tfam, NRF-1 and NRF-2, in rat liver, testis and brain, 965
metabolon – Hypothesis: Hyperstructures regulate bacterial structure and the cell cycle, 915
metamorphosis – Amphibian metamorphosis as a model for studying the developmental actions of thyroid hormone, 359
methyltransferase – *mraW*, an essential gene at the *dcw* cluster of *Escherichia coli* codes for a cytoplasmic protein with methyltransferase activity, 879
mitochondria – Oxidative DNA damage processing in nuclear and mitochondrial DNA, 155 – Analysis of wheat mitochondrial complex I purified by a one-step immunoaffinity chromatography, 645 – Cloning and characterisation of a pea mitochondrial NDPK, 1089 – Aging, mitochondria, and coenzyme Q_{10}: The neglected relationship, 1131
mitochondrial gene expression – Expression of mitochondrial genes and of the transcription factors involved in the biogenesis of mitochondria Tfam, NRF-1 and NRF-2, in rat liver, testis and brain, 965
mitochondrial transcription factor A – Expression of mitochondrial genes and of the transcription factors involved in the biogenesis of mitochondria Tfam, NRF-1 and NRF-2, in rat liver, testis and brain, 965
mitosis – DNA structure checkpoint pathways in *Schizosaccharomyces pombe*, 173
molecular combing – Single molecule analysis of DNA replication, 859
molecular evolution – Aminoacyl-tRNA synthetases: A new image for a classical family, 683
molecular genetics – Mutations in the sodium/iodide symporter (NIS) gene as a cause for iodide transport defects and congenital hypothyroidism, 469
molecular matchmaker – Nucleotide excision repair: From *E. coli* to man, 15
molten globule – Formation of a molten-globule-like state of cytochrome *c* induced by high concentrations of glycerol, 745
mouse model – Cancer from the outside, aging from the inside: Mouse models to study the consequences of defective nucleotide excision repair, 127
MraW protein – *mraW*, an essential gene at the *dcw* cluster of *Escherichia coli* codes for a cytoplasmic protein with methyltransferase activity, 879
multinodular goiter – Combined ultrasonographic and cytological studies in the diagnosis of thyroid nodules, 447
mutagenesis – Repair and mutagenesis survey of 8-hydroxyguanine in bacteria and human cells, 147 – Isoleucine 10 is essential for DNA gyrase B function in *Escherichia coli*, 973 – Display of Ras on filamentous phage through cysteine replacement, 1079
mutagenesis and carcinogenesis – Base excision repair of 8-hydroxyguanine protects DNA from endogenous oxidative stress, 59
mutant TR – DNA binding affinity of hTRβ1 mutants as heterodimers with traps from different tissues, 297
mutation – Mutation of the Secys residue 266 in human type 2 selenodeiodinase alters ^{75}Se incorporation without affecting its biochemical properties, 535
mycoplasma – Cell reproduction cycle of mycoplasma, 873
***myo*-inositol** – *myo*-[^{3}H]-inositol loaded erythrocytes and white ghosts: Two models to investigate the phosphatidylinositol synthesis in human red cells, 1011

N

NADPH oxidase – Biochemical characterization of a Ca^{2+}/NAD(P)H-dependent H_2O_2 generator in human thyroid tissue, 373
NF-Y – Analysis of the human HLA-DRA gene upstream region: Evidence for a stem-loop array directed by nuclear factors, 219
Nijmegen Breakage Syndrome – Mammalian X-ray-sensitive mutants which are defective in non-homologous (illegitimate) DNA double-strand break repair, 107
NKX2.1 – Thyroid-specific enhancer-binding protein/thyroid transcription factor 1 is not required for the initial specification of the thyroid and lung primordia, 321
Nod factors – Recent advances in the study of Nod factor perception and signal transduction, 669
non-equilibrium structure – Hypothesis: Hyperstructures regulate bacterial structure and the cell cycle, 915
nuclear proteins – Effect of phosphorothioate modifications on the ability of GTn oligodeoxynucleotides to specifically recognize single-stranded DNA-binding proteins and to affect human cancer cellular growth, 1115
nuclear receptors – Evolving concepts of thyroid hormone action, 539
nuclear respiratory factors 1 and 2 – Expression of mitochondrial genes and of the transcription factors involved in the biogenesis of mitochondria Tfam, NRF-1 and NRF-2, in rat liver, testis and brain, 965
nucleoids – Do plastid envelope membranes play a role in the expression of the plastid genome?, 619
nucleoside diphosphate kinase – Cloning and characterisation of a pea mitochondrial NDPK, 1089
nucleosome – The site of binding of linker histone to the nucleosome does not depend upon the amino termini of core histones, 727
nucleotide excision repair – Chromatin rearrangements during nucleotide excision repair, 45
nucleotide excision repair – Repair and mutagenesis survey of 8-hydroxyguanine in bacteria and human cells, 147
NusA – Nus A is involved in transcriptional termination on λ *tI*, 757
NusB – Nus A is involved in transcriptional termination on λ *tI*, 757

O–P

oligomerization – RNA:pseudouridine synthetase Pus1 from *Saccharomyces cerevisiae*: Oligomerization property and stoichiometry of the complex with yeast $tRNA^{Phe}$, 751 – Functional domains of DnaA proteins, 819
o-phthalaldehyde – Identification of lysine residue involved in inactivation of brain glutamate dehydrogenase isoproteins by o-phthalaldehyde, 1123
oriC – Cellular localization of *oriC* during the cell cycle of *Escherichia coli* as analyzed by fluorescent in situ hybridization, 797 – Chromosomal insertions localized around *oriC* affect the cell cycle in *Escherichia coli*, 811 – Initiation of DNA replication in Δ*cya* mutants of *Escherichia coli* K12, 827 – Replication cycle dependent association of SeqA to the outer membrane fraction of *E. coli*, 841
oxidative DNA damage – Oxidative DNA damage processing in nuclear and mitochondrial DNA, 155

6-phosphogluconate – Characterization of *Aspergillus niger* phosphoglucose isomerase. Use for quantitative determination of erythrose 4-phosphate, 267
p75^{hNTR} – Mechanisms of apical protein sorting in polarized thyroid epithelial cells, 347
papillary – Different mutations of the RET gene cause different human tumoral diseases, 397
peptide mapping – Identification of lysine residue involved in inactivation of brain glutamate dehydrogenase isoproteins by o-phthalaldehyde, 1123
perception and signal transduction – Recent advances in the study of Nod factor perception and signal transduction, 669
peroxidase – Molecular evolution of thyroid peroxidase, 557
pertussis toxin – An adenosine receptor agonist-induced modulation of TSH-dependent cell growth in FRTL-5 thyroid cells mediated by inhibitory G protein, G_i, 341
phage display – Display of Ras on filamentous phage through cysteine replacement, 1079
phosphoglucose isomerase – Characterization of *Aspergillus niger* phosphoglucose isomerase. Use for quantitative determination of erythrose 4-phosphate, 267
phosphoinositides – *myo*-[^{3}H]-inositol loaded erythrocytes and white ghosts: Two models to investigate the phosphatidylinositol synthesis in human red cells, 1011
phosphokinase – Proteoglycans in human malignant mesothelioma. Stimulation of their synthesis induced by epidermal, insulin and platelet-derived growth factors involves receptors with tyrosine kinase activity, 733
phospholipase A_2 – Signaling pathways involved in thyroid hyperfunction and growth in Graves' disease, 415
phosphorothioate oligonucleotides – Effect of phosphorothioate modifications on the ability of GTn oligodeoxynucleotides to specifically recognize single-stranded DNA-binding proteins and to affect human cancer cellular growth, 1115
photo-cross-links – A Y form of hammerhead ribozyme trapped by photo-cross-links retains full cleavage activity, 201
photoreactivation – The discovery of enzymatic photoreactivation and the question of priority: The letters of Salvador Luria and Albert Kelner, 7
Pisum sativum – Cloning and characterisation of a pea mitochondrial NDPK, 1089
plant – Protein retention and localization in the endoplasmic reticulum and the Golgi apparatus, 607
plant Golgi – Membrane trafficking in higher plant cells: GFP and antibodies, partners for probing the secretory pathway, 597
plant terminal membranes – Recent advances on proteins of plant terminal membranes, 577
plasma membrane – Involvement of plasma membrane proteins in plant defense responses. Analysis of the cryptogein signal transduction in tobacco, 663 – Large scale characterization of plant plasma membrane proteins, 655
plasminogen activation – Urokinase-type plasminogen activator and normal thyroid cell adhesion to the extracellular matrix, 355
plastid envelope – Do plastid envelope membranes play a role in the expression of the plastid genome?, 619
PNP oxidase – Refolding of pyridoxine-5'-P oxidase assisted by GRoeL, 1057
pollen-pistil interaction – Membrane proteins involved in pollen-pistil interactions, 675
poly(ADP-ribose) polymerase – Involvement of poly(ADP-ribose) polymerase in base excision repair, 69
postembryonic development – Amphibian metamorphosis as a model for studying the developmental actions of thyroid hormone, 359
post-transcription regulation – Acylphosphatase expression during macrophage differentiation and activation of U-937 cell line, 1031
preterm and term newborns – Hormone synthesis and storage in the thyroid of human preterm and term newborns: effect of thyroxine treatment, 563
PriA protein – *Escherichia coli* and *Bacillus subtilis* PriA proteins essential for recombination-dependent DNA replication: Involvement of ATPase/helicase activity of PriA for inducible stable DNA replication, 847
proliferation – Identification and characterization of mRNAs differentially expressed in thyroid cells stimulated by a mitogenic treatment, 309
promoter – Analysis of the human HLA-DRA gene upstream region: Evidence for a stem-loop array directed by nuclear factors, 219
protein catabolism – Distribution and subcellular localization of acylpeptide hydrolase and acylase I along the hog gastrointestinal tract, 1049
protein chemistry and structure – Sarcolectin: Complete purification for molecular cloning, 701
protein import – Cloning and characterisation of a pea mitochondrial NDPK, 1089
protein kinase – Regulation of the DNA-dependent protein kinase (DNA-PK) activity in eucaryotic cells, 117
protein synthesis – Aminoacyl-tRNA synthetases: A new image for a classical family, 683
protein synthesis – Identification of a ribosomal ATPase in *Escherichia coli* cells, 1097
proteins – Recent advances on proteins of plant terminal membranes, 577
proteoglycans – Purification and characterisation of a minor low-sulphated dermatan sulphate-proteoglycan from ray skin, 187 – Proteoglycans in human malignant mesothelioma. Stimulation of their synthesis induced by epidermal, insulin and platelet-derived growth factors involves receptors with tyrosine kinase activity, 733
proteome – Large scale characterization of plant plasma membrane proteins, 655
protooncogenes – Identification and characterization of mRNAs differentially expressed in thyroid cells stimulated by a mitogenic treatment, 309
Pseudomonas fragi – Acyl-CoA synthetase catalyzes the synthesis of diadenosine hexaphosphate (Ap_6A), 229
pseudouridine synthetase – RNA:pseudouridine synthetase Pus1 from *Saccharomyces cerevisiae*: Oligomerization property and stoichiometry of the complex with yeast $tRNA^{Phe}$, 751

R

RAD51 pathway – The contribution of homologous recombination in preserving genome integrity in mammalian cells, 87
radioiodine – Serum and tissue thyroglobulin measurement: Clinical applications in thyroid disease, 463
Raf1 – Display of Ras on filamentous phage through cysteine replacement, 1079
Ras – Display of Ras on filamentous phage through cysteine replacement, 1079
rat – Effects of iodine repletion on thyroid morphology in iodine and/or selenium deficient rat term fetuses, pups and mothers, 485
rat tissues – Expression of mitochondrial genes and of the transcription factors involved in the biogenesis of mitochondria Tfam, NRF-1 and NRF-2, in rat liver, testis and brain, 965
rate of cell division – Visualizing multiple constrictions in spheroidal *Escherichia coli* cells, 897
ray skin – Purification and characterisation of a minor low-sulphated dermatan sulphate-proteoglycan from ray skin, 187
reactive lysine – Identification of lysine residue involved in inactivation of brain glutamate dehydrogenase isoproteins by o-phthalaldehyde, 1123
receptor-like kinases – Membrane proteins involved in pollen-pistil interactions, 675
recombinant DNA – Sarcolectin (SCL): Structure and expression of the recombinant molecule, 709
recombinational repair – The contribution of homologous recombination in preserving genome integrity in mammalian cells, 87
refolding process – Refolding of pyridoxine-5'-P oxidase assisted by GRoeL, 1057
regulation – Regulation of the DNA-dependent protein kinase (DNA-PK) activity in eucaryotic cells, 117 – Radiation inducible

DNA repair processes in eukaryotes, 161
replication – Chromosomal insertions localized around *oriC* affect the cell cycle in *Escherichia coli*, 811 – Initiation of DNA replication in Δ*cya* mutants of *Escherichia coli* K12, 827
replisome – *Escherichia coli* contains a DNA replication compartment in the cell center, 803
resistance – Involvement of plasma membrane proteins in plant defense responses. Analysis of the cryptogein signal transduction in tobacco, 663
restriction enzyme – Post-segregational killing by restriction modification gene complexes: Observations of individual cell deaths, 931
retention signal – Protein retention and localization in the endoplasmic reticulum and the Golgi apparatus, 607
ribosomal protein – In vitro assembly of yeast 5S rRNA and a fusion protein containing ribosomal protein L5 and maltose binding protein, 1015
ribosomes – Identification of a ribosomal ATPase in *Escherichia coli* cells, 1097
RNA polymerase – Effect of DNA lesions on transcription elongation, 139
RNA polymerase binding – Detection of an extended –10 element in the promoter region of the *pckA* gene encoding phosphoenolpyruvate carboxykinase in *Escherichia coli*, 197
RNA recognition – Aminoacyl-tRNA synthetases: A new image for a classical family, 683
RNA-protein interaction – In vitro assembly of yeast 5S rRNA and a fusion protein containing ribosomal protein L5 and maltose binding protein, 1015
RXRs – DNA binding affinity of hTRβ1 mutants as heterodimers with traps from different tissues, 297

S

^{75}Se incorporation – Mutation of the Secys residue 266 in human type 2 selenodeiodinase alters ^{75}Se incorporation without affecting its biochemical properties, 535
5S rRNA – In vitro assembly of yeast 5S rRNA and a fusion protein containing ribosomal protein L5 and maltose binding protein, 1015
***S* RNases** – Membrane proteins involved in pollen-pistil interactions, 675
S. cerevisiae – Radiation inducible DNA repair processes in eukaryotes, 161
S_1 mapping – Nus A is involved in transcriptional termination on λ *tI*, 757
Salvador Luria – The discovery of enzymatic photoreactivation and the question of priority: The letters of Salvador Luria and Albert Kelner, 7
sarcolectin – Sarcolectin: Complete purification for molecular cloning, 701
Schizosaccharomyces pombe – DNA structure checkpoint pathways in *Schizosaccharomyces pombe,* 173
SCL – Sarcolectin (SCL): Structure and expression of the recombinant molecule, 709
SECIS binding protein – A SECIS binding protein (SBP) is distinct from selenocystyl-tRNA protecting factor (SePF), 213
SECIS RNA – A SECIS binding protein (SBP) is distinct from selenocystyl-tRNA protecting factor (SePF), 213
secondary structures – The structures of elastins and their function, 981
Sec-tRNA^{Sec} protecting factor – A SECIS binding protein (SBP) is distinct from selenocystyl-tRNA protecting factor (SePF), 213
selenium – Effects of iodine repletion on thyroid morphology in iodine and/or selenium deficient rat term fetuses, pups and mothers, 485 – The trace element selenium and the thyroid gland, 527
selenocysteine – A SECIS binding protein (SBP) is distinct from selenocystyl-tRNA protecting factor (SePF), 213
selenodeiodinase – Mutation of the Secys residue 266 in human type 2 selenodeiodinase alters ^{75}Se incorporation without affecting its biochemical properties, 535
selenoprotein – The trace element selenium and the thyroid gland, 527
self-association – Annexin VI interacts with adenine nucleotides and their analogs, 717
self-incompatibility – Membrane proteins involved in pollen-pistil interactions, 675
selfish gene – Post-segregational killing by restriction modification gene complexes: Observations of individual cell deaths, 931
seqA protein – Replication cycle dependent association of SeqA to the outer membrane fraction of *E. coli*, 841
sequence – Sarcolectin (SCL): Structure and expression of the recombinant molecule, 709
serum lipoproteins – Altered thyroid hormone binding to plasma lipoproteins in the syndrome of resistance to thyroid hormones, 545
Shine-Dalgarno – A symbolic-numeric approach to find patterns in genomes. Application to the translation initiation sites of *E. coli*, 1065
signal transduction – Involvement of plasma membrane proteins in plant defense responses. Analysis of the cryptogein signal transduction in tobacco, 663
skin fibroblasts – GSH system in relation to redox state in dystrophic skin fibroblasts, 1025
***S*-locus** – Membrane proteins involved in pollen-pistil interactions, 675
solid phase translation – Initiation of *Escherichia coli* ribosomes on matrix coupled mRNAs studied by optical biosensor technique, 995
soluble proteins – Protein retention and localization in the endoplasmic reticulum and the Golgi apparatus, 607
somatomedin – Endogenous insulin-like growth factors regulate the proliferation of TSH-independent mutants derived from FRTL5 cells, 367
splenocyte – Upregulation of glyceraldehyde-3-phosphate dehydrogenase mRNA in the spleen of tumor-bearing mice, 1109
Streptomyces – Functional domains of DnaA proteins, 819
structural disorder – Disordered C-terminal domain of tyrosyl-tRNA synthetase: Secondary structure prediction, 235
structural prediction – Disordered C-terminal domain of tyrosyl-tRNA synthetase: Secondary structure prediction, 235
subcellular fractionation – Distribution and subcellular localization of acylpeptide hydrolase and acylase I along the hog gastro-intestinal tract, 1049
substrate properties – Low enantioselectivities of human deoxycytidine kinase and human deoxyguanosine kinase with respect to 2'-deoxyadenosine, 2'-deoxyguanosine and their analogs, 1041
substrate specificity – Comparative studies of Ca^{2+}-dependent proteases (CDP I and CDP II) from *Allomyces arbuscula*, 765
synchronous culture – Division-associated changes in membrane viscosity of *Escherichia coli*, 921

T

***T/ebp*-null mouse** – Thyroid-specific enhancer-binding protein/thyroid transcription factor 1 is not required for the initial specification of the thyroid and lung primordia, 321
TFIIH – Transcription factor IIH: A key player in the cellular response to DNA damage, 27
Thermus – Functional domains of DnaA proteins, 819
thiouracil – Summary of observations on transplantable tumors of the rat thyroid gland, 409
threading – Disordered C-terminal domain of tyrosyl-tRNA synthetase: Secondary structure prediction, 235
thylakoid – Mutants of *Chlamydomonas*: Tools to study thylakoid membrane structure, function and biogenesis, 631
thymine limitation – Visualizing multiple constrictions in spheroidal *Escherichia coli* cells, 897
thyrocyte – Biochemical characterization of a Ca^{2+}/NAD(P)H-dependent H_2O_2 generator in human thyroid tissue, 373
thyroglobulin – Thyroglobulin regulates follicular function and heterogeneity by suppressing thyroid-specific gene expression, 329 – The screening for mutations in the thyroglobulin cDNA from six patients with congenital hypothyroidism,

425 – Colloidal aggregates of insoluble inclusions in human goiters, 441 – Serum and tissue thyroglobulin measurement: Clinical applications in thyroid disease, 463 – Depletion of divalent cations within the secretory pathway inhibits the terminal glycosylation of complex carbohydrates of thyroglobulin, 497 – The importance of thyroglobulin structure for thyroid hormone biosynthesis, 505 – The disulfide bond pattern between fragments obtained by the limited proteolysis of bovine thyroglobulin, 517 – Selective endocytosis of thyroglobulin: A review of potential mechanisms for protecting newly synthesized molecules from premature degradation, 549 – Hormone synthesis and storage in the thyroid of human preterm and term newborns: effect of thyroxine treatment, 563

thyroglobulin receptor – Thyroglobulin binding and TSH regulation of the RHL-1 subunit of the asialoglycoprotein receptor in rat thyroid, 493 – Depletion of divalent cations within the secretory pathway inhibits the terminal glycosylation of complex carbohydrates of thyroglobulin, 497

thyroid – Identification and characterization of mRNAs differentially expressed in thyroid cells stimulated by a mitogenic treatment, 309 – Urokinase-type plasminogen activator and normal thyroid cell adhesion to the extracellular matrix, 355 – Endogenous insulin-like growth factors regulate the proliferation of TSH-independent mutants derived from FRTL5 cells, 367 – Biochemical characterization of a Ca^{2+}/NAD(P)H-dependent H_2O_2 generator in human thyroid tissue, 373 – Different mutations of the RET gene cause different human tumoral diseases, 397 – The screening for mutations in the thyroglobulin cDNA from six patients with congenital hypothyroidism, 425 – Cloning, chromosomal localization and identification of polymorphisms in the human thyroid transcription factor 2 gene (TITF2), 433 – Colloidal aggregates of insoluble inclusions in human goiters, 441 – Mutations in the sodium/iodide symporter (NIS) gene as a cause for iodide transport defects and congenital hypothyroidism, 469 – Effects of iodine repletion on thyroid morphology in iodine and/or selenium deficient rat term fetuses, pups and mothers, 485 – The trace element selenium and the thyroid gland, 527

thyroid autoimmunity – Signaling pathways involved in thyroid hyperfunction and growth in Graves' disease, 415

thyroid cancer – Thyroid-specific gene expression in the multi-step process of thyroid carcinogenesis, 389 – Combined ultrasonographic and cytological studies in the diagnosis of thyroid nodules, 447 – Serum and tissue thyroglobulin measurement: Clinical applications in thyroid disease, 463

thyroid cells – Response to thyrotropin of normal thyroid follicular cell strain FRTL5 in hypergravity, 281 – Effects of thyroid-stimulating hormone on the replicative cycle of vesicular stomatitis virus in primary cultured sheep thyroid cells, 291 – An adenosine receptor agonist-induced modulation of TSH-dependent cell growth in FRTL-5 thyroid cells mediated by inhibitory G protein, G_i, 341 – Thyroid-specific gene expression in the multi-step process of thyroid carcinogenesis, 389 – Expression of integrins of the β family in thyroid cells from patients with Graves' disease in vivo and in vitro, 477

thyroid differentiation – Thyroglobulin binding and TSH regulation of the RHL-1 subunit of the asialoglycoprotein receptor in rat thyroid, 493

thyroid follicles – Induction of apoptosis in porcine thyroid follicles by transforming growth factor β1 and epidermal growth factor, 315

thyroid gland – Hormone synthesis and storage in the thyroid of human preterm and term newborns: effect of thyroxine treatment, 563

thyroid hormone – Amphibian metamorphosis as a model for studying the developmental actions of thyroid hormone, 359

thyroid hormone action – Evolving concepts of thyroid hormone action, 539

thyroid hormone binding – Altered thyroid hormone binding to plasma lipoproteins in the syndrome of resistance to thyroid hormones, 545

thyroid hormone metabolism – Tissue-specific patterns of changes in 3,5,3´-triiodo-L-thyronine concentrations in thyroidectomized rats infused with increasing doses of the hormone. Which are the regulatory mechanisms?, 453

thyroid hormone receptor – DNA binding affinity of hTRβ1 mutants as heterodimers with traps from different tissues, 297 – Amphibian metamorphosis as a model for studying the developmental actions of thyroid hormone, 359

thyroid hormone resistance – Altered thyroid hormone binding to plasma lipoproteins in the syndrome of resistance to thyroid hormones, 545

thyroid hormone synthesis – Selective endocytosis of thyroglobulin: A review of potential mechanisms for protecting newly synthesized molecules from premature degradation, 549

thyroid hormones – The importance of thyroglobulin structure for thyroid hormone biosynthesis, 505

thyroid morphogenesis – Thyroid-specific enhancer-binding protein/thyroid transcription factor 1 is not required for the initial specification of the thyroid and lung primordia, 321

thyroid nodule – Combined ultrasonographic and cytological studies in the diagnosis of thyroid nodules, 447

thyroid peroxidase – Molecular evolution of thyroid peroxidase, 557

thyroid polarity – Mechanisms of apical protein sorting in polarized thyroid epithelial cells, 347

thyroid stimulating hormone – Effects of thyroid-stimulating hormone on the replicative cycle of vesicular stomatitis virus in primary cultured sheep thyroid cells, 291

thyroid transcription factor 1 (TTF1) – Thyroid-specific enhancer-binding protein/thyroid transcription factor 1 is not required for the initial specification of the thyroid and lung primordia, 321

thyroid transcription factors – Thyroid-specific gene expression in the multi-step process of thyroid carcinogenesis, 389

thyroid-specific enhancer-binding protein (T/EBP) – Thyroid-specific enhancer-binding protein/thyroid transcription factor 1 is not required for the initial specification of the thyroid and lung primordia, 321

thyroid-specific gene expression – Thyroglobulin regulates follicular function and heterogeneity by suppressing thyroid-specific gene expression, 329

thyrotropin – Response to thyrotropin of normal thyroid follicular cell strain FRTL5 in hypergravity, 281 – Inhibition of farnesylation blocks growth but not differentiation in FRTL-5 thyroid cells, 287 – Induction of apoptosis in porcine thyroid follicles by transforming growth factor β1 and epidermal growth factor, 315 – An adenosine receptor agonist-induced modulation of TSH-dependent cell growth in FRTL-5 thyroid cells mediated by inhibitory G protein, G_i, 341

thyroxine – Biochemical characterization of a Ca^{2+}/NAD(P)H-dependent H_2O_2 generator in human thyroid tissue, 373 – Tissue-specific patterns of changes in 3,5,3´-triiodo-L-thyronine concentrations in thyroidectomized rats infused with increasing doses of the hormone. Which are the regulatory mechanisms?, 453 – Hormone synthesis and storage in the thyroid of human preterm and term newborns: effect of thyroxine treatment, 563

tobacco – Involvement of plasma membrane proteins in plant defense responses. Analysis of the cryptogein signal transduction in tobacco, 663

topoisomerase II – The catalytic activities of DNA topoisomerase II are most closely associated with the DNA cleavage/religation steps, 771

topoisomerase inhibitors – The catalytic activities of DNA topoisomerase II are most closely associated with the DNA cleavage/religation steps, 771

TPO – Molecular evolution of thyroid peroxi-

dase, 557
transcription – Transcription factor IIH: A key player in the cellular response to DNA damage, 27
transcription factor – Cloning, chromosomal localization and identification of polymorphisms in the human thyroid transcription factor 2 gene (TITF2), 433
transcription termination – Nus A is involved in transcriptional termination on λ *tI*, 757
transcriptional regulation – Control of galectin gene expression, 381
transcription-coupled repair – Effect of DNA lesions on transcription elongation, 139
transfer – Tissue-specific patterns of changes in 3,5,3´-triiodo-L-thyronine concentrations in thyroidectomized rats infused with increasing doses of the hormone. Which are the regulatory mechanisms?, 453
transforming growth factor-β1 – Induction of apoptosis in porcine thyroid follicles by transforming growth factor β1 and epidermal growth factor, 315
translation – Do plastid envelope membranes play a role in the expression of the plastid genome?, 619
translation factors – Identification of a ribosomal ATPase in *Escherichia coli* cells, 1097
translation initiation – A symbolic-numeric approach to find patterns in genomes. Application to the translation initiation sites of *E. coli*, 1065
TRAPs – DNA binding affinity of hTRβ1 mutants as heterodimers with traps from different tissues, 297
triiodothyronine – Biochemical characterization of a Ca^{2+}/NAD(P)H-dependent H_2O_2 generator in human thyroid tissue, 373
triiodothyronine – Tissue-specific patterns of changes in 3,5,3´-triiodo-L-thyronine concentrations in thyroidectomized rats infused with increasing doses of the hormone. Which are the regulatory mechanisms?, 453 – Evolving concepts of thyroid hormone action, 539
tRNA – RNA:pseudouridine synthetase Pus1 from *Saccharomyces cerevisiae*: Oligomerization property and stoichiometry of the complex with yeast $tRNA^{Phe}$, 751
tRNA identity – Aminoacyl-tRNA synthetases: A new image for a classical family, 683
TSH receptor – Signaling pathways involved in thyroid hyperfunction and growth in Graves' disease, 415
TSH regulation – Thyroglobulin binding and TSH regulation of the RHL-1 subunit of the asialoglycoprotein receptor in rat thyroid, 493
tumor marker – Serum and tissue thyroglobulin measurement: Clinical applications in thyroid disease, 463
tumor progression – Control of galectin gene expression, 381 – Summary of observations on transplantable tumors of the rat thyroid gland, 409
two-dimensional gel electrophoresis – Large scale characterization of plant plasma membrane proteins, 655
tyrosine – Different mutations of the RET gene cause different human tumoral diseases, 397
tyrosyl-tRNA synthetase – Disordered C-terminal domain of tyrosyl-tRNA synthetase: Secondary structure prediction, 235

U

U-937 – Acylphosphatase expression during macrophage differentiation and activation of U-937 cell line, 1031
ubiquinone – Aging, mitochondria, and coenzyme Q_{10}: The neglected relationship, 1131
UGA codon – A SECIS binding protein (SBP) is distinct from selenocystyl-tRNA protecting factor (SePF), 213
ultrasonography – Combined ultrasonographic and cytological studies in the diagnosis of thyroid nodules, 447
ultraviolet light – DNA damage recognition during nucleotide excision repair in mammalian cells, 39 – Effect of DNA lesions on transcription elongation, 139
ultraviolet radiation – The discovery of enzymatic photoreactivation and the question of priority: The letters of Salvador Luria and Albert Kelner, 7
urokinase – Urokinase-type plasminogen activator and normal thyroid cell adhesion to the extracellular matrix, 355
urokinase receptor – Urokinase-type plasminogen activator and normal thyroid cell adhesion to the extracellular matrix, 355

V–W

V(D)J recombination – Mammalian X-ray-sensitive mutants which are defective in non-homologous (illegitimate) DNA double-strand break repair, 107
vesicle trafficking – Recent advances on proteins of plant terminal membranes, 577
vesicular stomatitis virus – Effects of thyroid-stimulating hormone on the replicative cycle of vesicular stomatitis virus in primary cultured sheep thyroid cells, 291
water – Hypothesis: Hyperstructures regulate bacterial structure and the cell cycle, 915
Western blots – Comparative studies of Ca^{2+}-dependent proteases (CDP I and CDP II) from *Allomyces arbuscula*, 765
wheat – Analysis of wheat mitochondrial complex I purified by a one-step immunoaffinity chromatography, 645

X–Y

xeroderma pigmentosum – DNA damage recognition during nucleotide excision repair in mammalian cells, 39
X-ray-sensitive mutants – Mammalian X-ray-sensitive mutants which are defective in non-homologous (illegitimate) DNA double-strand break repair, 107
yeast – In vitro assembly of yeast 5S rRNA and a fusion protein containing ribosomal protein L5 and maltose binding protein, 1015